

S2 Table. Prevalence of currently receiving treatment of selected diseases of the study population

	Men (N=27,717)	Women (N=56,377)	<i>P</i> ^a
<i>Certain infectious & parasitic disease</i>			
Tuberculosis, <i>n</i> (%)	23 (0.1)	28 (0.1)	< 0.001
<i>Viral infections characterized by skin & mucous membrane lesions</i>			
Acute liver disease, <i>n</i> (%)	44 (0.2)	28 (0.1)	< 0.001
Chronic liver disease, <i>n</i> (%)	273 (1.0)	242 (0.4)	< 0.001
<i>Neoplasms</i>			
Cancer, <i>n</i> (%)	140 (0.5)	409 (0.7)	< 0.001
<i>Endocrine, nutritional & metabolic disease</i>			
Diabetes mellitus, <i>n</i> (%)	1,966 (7.1)	2,402 (4.3)	< 0.001
Thyroid disease, <i>n</i> (%)	143 (0.5)	1,228 (2.2)	< 0.001
Hyperlipidemia, <i>n</i> (%)	737 (2.7)	1,560 (2.7)	0.130
<i>Mental & behavioral disorders</i>			
Depression, <i>n</i> (%)	108 (0.4)	461 (0.8)	< 0.001
<i>Diseases of the eye & adnexa</i>			
Cataract, <i>n</i> (%)	185 (0.7)	375 (0.7)	< 0.001
<i>Diseases of the circulatory system</i>			
Hypertension, <i>n</i> (%)	5,033 (18.2)	8,685 (15.4)	< 0.001
Myocardial infarction, <i>n</i> (%)	564 (2.0)	574 (1.0)	< 0.001
Stroke, <i>n</i> (%)	273 (1.0)	268 (0.5)	< 0.001
<i>Diseases of the respiratory system</i>			
Asthma / Chronic bronchitis, <i>n</i> (%)	161 (0.6)	318 (0.6)	0.616
<i>Diseases of the digestive system</i>			
Cholelithiasis, <i>n</i> (%)	34 (0.1)	63 (0.1)	0.888
Fatty liver disease, <i>n</i> (%)	302 (1.1)	222 (0.4)	< 0.001
Gastritis, <i>n</i> (%)	416 (1.5)	1,050 (1.9)	< 0.001
Intestinal polyp, <i>n</i> (%)	33 (0.1)	31 (0.1)	< 0.001
Peptic Ulcer, <i>n</i> (%)	285 (1.0)	464 (0.8)	0.012
<i>Diseases of the musculo-skeletal system & connective tissue</i>			
Arthritis, <i>n</i> (%)	350 (1.3)	2,871 (5.1)	< 0.001
Osteoporosis, <i>n</i> (%)	45 (0.2)	1,667 (3.0)	< 0.001
<i>Diseases of the genitourinary system</i>			
Bladder infection, <i>n</i> (%)	33 (0.1)	225 (0.4)	< 0.001

a. *p*-value was calculated by chi-square test for categorical variable