

Supplemental Table. Clinical characteristics of the dry eye study population and galectin-3 levels.

Patient No	Sex	Age	Eye	ATD	MGD	GVHD	SS	Gal-3 in tears (ng/μl)	Gal-3C/Gal-3 Ratio
1	F	50	OD	+	-	-	-	0.27	0.00
2	F	53	OD	+	-	-	-	0.55	0.00
3	F	58	OD	+	-	-	-	0.04	0.00
4	M	50	OD	+	-	-	-	1.33	0.29
5	F	54	OD	+	-	-	-	0.41	0.39
6	F	62	OD	+	-	-	-	0.09	0.00
7	F	56	OD	+	-	-	-	0.17	0.00
8	F	59	OD	+	-	-	-	0.30	0.36
9	M	52	OD	+	-	-	-	0.38	0.00
10	M	47	OD	+	-	-	-	0.14	0.00
11	F	63	OD	+	-	-	-	0.08	0.00
12	F	57	OD	+	-	-	-	1.36	0.23
13	M	57	OD	+	+	+	-	0.13	0.18
			OS	+	+	+	-	0.19	0.52
14	M	35	OD	+	+	+	-	0.35	0.39
			OS	+	+	+	-	0.32	0.42
15	F	84	OD	+	+	-	+	0.71	0.34
			OS	+	+	-	+	0.35	0.00
16	M	74	OD	-	+	-	-	0.09	0.00
			OS	-	+	-	-	0.32	0.02

ATD = aqueous tear deficiency; DE = dry eye; Gal-3 = galectin-3; Gal-3C = galectin-3C; GVHD = graft versus host disease; MGD = meibomian gland dysfunction; SS = Sjögren syndrome