

## Supplementary Online Content

Boyle CA, Perrin JM, Moyer VA. Use of clinical preventive services in infants, children, and adolescents. *JAMA*. doi:10.1001/jama.2014.12890.

**eTable.** Percentage of patients who are receiving selected recommended clinical preventive services for infants, children, and adolescents — United States

This supplementary material has been provided by the authors to give readers additional information about their work.

© 2014 American Medical Association. All rights reserved.

<b>eTABLE. Percentage of patients who are receiving selected recommended clinical preventive services for infants, children, and adolescents — United States</b>		
<b>Topic/Indicator (years received) and Data Sources</b>	<b>% receiving service</b>	<b>Clinical practice recommendation or guidelines for the preventive service – recommending organization(s)</b>
<b>Breastfeeding counseling (2010)</b> <b>Data source: Pregnancy Risk Assessment Monitoring System</b>		
Women with recent live births who reported receiving any advice about breastfeeding during prenatal care visits	82.7	USPSTF/ACOG/AAP
<b>Hearing screening and follow-up (2009–2010)</b> <b>Data source: CDC Early Hearing Detection and Intervention Hearing Screening and Follow-up Survey</b>		
Infants who failed their hearing screening and then received diagnostic testing needed to confirm hearing loss	50.3	USPSTF/AAP
<b>Developmental screening (2007)</b> <b>Data source: National Survey of Children’s Health</b>		
Children aged 10–47 months whose parents were asked by health-care providers to complete a formal screen for developmental delays during the preceding 12 months	21.1	AAP
<b>Lead screening (2010)</b> <b>Data source: CDC Child Blood Lead Surveillance System</b>		
Children aged 1–2 years who were screened and reported to CDC for lead poisoning	33.4	AAP
<b>Vision screening (2009–2010)</b> <b>Data source: Medical Expenditure Panel Survey</b>		
Children aged 5 years who were reported by their parents to have ever had their vision checked by a doctor or other health provider	77.9	USPSTF/AA
<b>Hypertension screening (2009–2010)</b> <b>Data sources: National Ambulatory Medical Care Survey, National Hospital Ambulatory Medical Care Survey, and Medical Expenditure Panel Survey</b>		
Provider reported office-based and hospital outpatient department preventive care visits with documentation of blood pressure measurement among children and adolescents aged 3–17 years	75.7	AAP
Children and adolescents aged 3–17 years were reported by their parents or caregivers to have had their blood pressure measured by a doctor or other health provider at a nonemergency care physician or clinic visit during the preceding year	69.6	AAP
<b>Dental care and dental preventive services (2005–2010)</b> <b>Data sources: Medical Expenditure Panel Survey and National Health and Nutrition Examination Survey</b>		
Persons aged ≤21 years who have visited the dentist during the preceding year (2009)	43.8	AAP
Persons aged ≤21 years who have received dental preventive services (topical fluoride, sealant, or both) during the preceding year (2009)	14.2	AAP
Persons aged 5–19 years who have a dental sealant (2005–2010)	31.3	AAP
<b>Human papillomavirus (HPV) vaccination (2011)</b> <b>Data source: National Immunization Survey – Teen</b>		
Adolescents females aged 13–17 years who have received ≥1 dose of the HPV vaccine	53.0	ACIP
Adolescent females aged 13–17 years who have received ≥3 doses of the HPV vaccine	34.8	ACIP
Adolescent males aged 13–17 years who have received ≥1 dose of the HPV vaccine	8.3	ACIP
Adolescent males aged 13–17 years who have received ≥3 doses of the HPV vaccine	1.3	ACIP

Topic/Indicator (years received) and Data Sources	% receiving service	Clinical practice recommendation or guidelines for the preventive service – recommending organization(s)
<b>Tobacco use screening and cessation assistance (2004–2010)</b>		
<b>Data source: National Ambulatory Medical Care Survey</b>		
Provider reported office-based outpatient visits with documentation of tobacco use status among persons aged 11–21 years	69.5	USPHS/AAP
Provider reported office-based outpatient visits with documentation of tobacco cessation assistance, including counseling and/or a prescription or order for a cessation medication among current tobacco users in persons aged 11–21 years	19.8	USPHS/AAP
<b>Chlamydia screening (2005–2010)</b>		
<b>Data sources: National Survey of Family Growth and National Ambulatory Medical Care Survey</b>		
Sexually active females aged 15–21 years who reported being tested for chlamydia during the preceding 12 months (2006–2010)	40.0	USPSTF/GAPS/AAP
Provider reported office-based ambulatory care setting visits with screening for chlamydia among females aged 15–21 years (2005–2010)	4.3	USPSTF/GAPS/AAP
<b>Reproductive health services (2006–2010)</b>		
<b>Data source: National Survey of Family Growth</b>		
Sexually experienced females aged 15–19 years who reported receiving a reproductive health service from a health-care provider during the preceding 12 months	76.5	GAPS/AAP
All females aged 15–19 years who reported receiving a reproductive health service from a health-care provider during the preceding 12 months	43.9	GAPS/AAP
Sexually experienced males aged 15–19 years who reported receiving a reproductive health service from a health-care provider during the preceding 12 months	62.5	GAPS/AAP
All males aged 15–19 years who reported receiving a reproductive health service from a health-care provider during the preceding 12 months	58.2	GAPS/AAP

USPSTF: U.S. Preventive Services Task Force; AGOG: American Congress of Obstetricians and Gynecologists; AAP: American Academy of Pediatrics; USPHS: U.S. Public Health Service; ACIP: Advisory Committee on Immunization Practices; GAPS: Guidelines for Adolescent Preventive Services

© 2014 American Medical Association. All rights reserved.