

Supplementary 2. Pharmacokinetic parameters of unincorporated CIGB-300.

Parameter	Group I (35 mg) N = 7	Group II (70 mg) N = 7	p (95% CI)
Tmax (h)**	0.17 ± 1.36	0.02 ± 0.23	0.01 (0.1; 2.0)
Cmax (µg/mL)*	9.7 ± 12.5	14.1 ± 13.9	0.85 (-17.0; 6.5)
λ (h ⁻¹)**	0.33 ± 0.39	0.50 ± 0.17	0.69 (-0.3; 0.4)
t _{1/2} (h)*	1.9 ± 0.7	1.3 ± 0.6	0.02 (0.1; 1.7)
AUC (µg.h/mL)*	17.7 ± 19.6	13.9 ± 13.2	0.16 (-6.5; 32.0)
Vd/F (L/Kg)**	0.2 ± 0.6	0.1 ± 0.6	0.31 (-0.5; 1.0)
CL/F (L/h.Kg)**	0.09 ± 0.46	0.11 ± 0.55	0.54 (-0.6; 0.6)
MRT (h)*	2.7 ± 1.1	1.7 ± 1.1	0.18-1.2; 2.2)

*Mean ± standard deviation **Median ± quartile range.

Abbreviation: CI = confidence interval for the differences

p: probability that the difference was > 0