

1 **Supplemental Data**

2 **Perfluorooctanoic acid (PFOA)-induced liver lesions in two strains of mice following developmental**
 3 **exposures: PPAR α is not required.**

4 Adam J. Filgo^{1,2}, Erin M. Quist^{2,5}, Mark J. Hoenerhoff³, Amy E. Brix⁶, Grace E. Kissling⁴, and Suzanne
 5 E. Fenton²

6
 7 ¹Curriculum in Toxicology, University of North Carolina, School of Medicine, Chapel Hill, NC, USA

8 ²National Toxicology Program Laboratory and ³Cellular and Molecular Pathology Branch, Division of the
 9 National Toxicology Program; ⁴Biostatistics Branch, Division of Intramural Research;
 10 National Institute of Environmental Health Sciences, National Institute of Health, RTP, NC, USA

11 ⁵Comparative Biomedical Sciences, College of Veterinary Medicine, North Carolina State University,
 12 Raleigh, NC, USA

13 ⁶Experimental Pathology Laboratories, Inc., RTP, NC, USA

14
 15 Supplemental Data Table 1. Percentage (%) of mice euthanized prior to 18 months (end of study) in
 16 the three strains utilized in these studies.

Mouse Strain	PFOA Dose (mg PFOA/kg body weight/day)							
	0	0.01	0.1	0.3	0.6	1.0	3.0	5.0
CD-1*	27.5	17.1	15.9	27.7	n/a	24.4	n/a	22.2
129/Sv WT	16.7	n/a	16.7	11.1	14.3	11.1	n/a	n/a
129/Sv PPAR α KO	25.0	n/a	23.1	16.7	n/a	0	18.2	n/a

17 n/a = not applicable; this dose was not used in the strain listed.

18 * Average for CD-1 in 8 separate studies ending at 18 months was 21.7% (Giknis and Clifford, 2010).

19