

S2 Table. Captured bat species in tropical mountain cloud forest fragments (TMCF) and shade coffee plantations (SCP). Guild bat classification is based on Rojas et al. 2012.

Species	Guild	TMCF	SCP	Total
Stenodermatinae				
<i>Artibeus jamaicensis</i>	Frugivore	41	48	89
<i>Artibeus lituratus</i>	Frugivore	2	11	13
<i>Dermanura tolteca</i>	Frugivore	23	9	32
<i>Centurio senex</i>	Frugivore	0	3	3
<i>Sturnira hodurensis</i>	Frugivore	329	95	424
<i>Sturnira parvidens</i>	Frugivore	63	12	75
Carollinae				
<i>Carollia sowelli</i>	Frugivore	47	17	64
Glossophaginae				
<i>Anoura geoffroyi</i>	Nectarivore	3	13	16
<i>Glossophaga soricina</i>	Nectarivore	0	6	6
<i>Glossophaga commissarisi</i>	Nectarivore	1	5	6
Desmodontinae				
<i>Desmodus rotundus</i>	Saguinivore	13	0	13
Verpertelionidae				
<i>Myotis keaysi</i>	Insectivore	6	2	8
Mormoopidae				
<i>Mormoops megalophylla</i>	Insectivore	15	2	17
<i>Pteronotus davyi</i>	Insectivore	0	1	1
<i>Pteronotus parnelli</i>	Insectivore	1	0	1
Total		544	224	768
Richness (S)		12	14	15

Rojas D, Vale Á, Ferrero V, Navarro L (2012) The role of frugivory in the diversification of bats in the Neotropics. *Journal of Biogeography* 39: 1948–1960. doi:10.1111/j.1365-2699.2012.02709.x.