

fMRI measurements of amygdala activation are confounded by stimulus correlated signal fluctuation in nearby veins draining distant brain regions

Roland N. Boubela^{abc†}, Klaudius Kalcher^{abc†}, Wolfgang Huf^{abc},
Eva-Maria Seidel^d, Birgit Derntl^e, Lukas Pezawas^f, Christian
Našel^g, Ewald Moser^{abh*}

^a*Center for Medical Physics and Biomedical Engineering, Medical University of Vienna,
Vienna, Austria*

^b*MR Centre of Excellence, Medical University of Vienna, Vienna, Austria*

^c*Department of Statistics and Probability Theory, Vienna University of Technology, Vienna,
Austria*

^d*Social, Cognitive and Affective Neuroscience Unit, Department of Basic Psychological
Research and Research Methods, Faculty of Psychology, University of Vienna, Vienna, Austria*

^e*Department of Psychiatry, Psychotherapy and Psychosomatics,
RWTH Aachen University, Aachen, Germany*

^f*Department of Psychiatry and Psychotherapy, Medical University of Vienna, Vienna, Austria*

^g*Department of Radiology, Tulln Hospital, Karl Landsteiner University of Health Sciences,
Tulln, Austria*

^h*Brain Behaviour Laboratory, Department of Psychiatry,
University of Pennsylvania Medical Center, Philadelphia, PA, USA*

[†]*Roland N. Boubela and Klaudius Kalcher contributed equally to this work.*

**Correspondence to ewald.moser@meduniwien.ac.at*

February 26, 2015

Supplementary Video Single-subject SWI and Faces-Forms Activation