

**Table S3 Characteristics of excluded full-text studies**

<b>Studies</b>	<b>Reasons for exclusion</b>
Mahood 1991 <sup>1</sup>	Cancer type unclear, 5-fluorouracil related stomatitis
Rocke 1993 <sup>2</sup>	Cancer type unclear, 5-fluorouracil related stomatitis
Cascinu 1994 <sup>3</sup>	Colon-rectum, pancreas, stomach, prostate cancers, 5-fluorouracil induced stomatitis
Nikoletti 2005 <sup>4</sup>	Colorectal cancer, 5-fluorouracil related stomatitis, plain ice versus flavoured ice,
Karagozoglu 2005 <sup>5</sup>	Epidermoid cancer, small cell cancer, adenocarcinoma, mesothelioma
Baydar 2005 <sup>6</sup>	Carcinomas of the colon, rectum, stomach, head of pancreas, cecum, sigmoid, neuroendocrine tumors, and metastatic tumors of unknown origin of the liver.
Papadeas 2007 <sup>7</sup>	Cancer type unclear
Katranci 2012 <sup>8</sup>	Gastric cancer, colon cancer, pancreatic cancer, and rectal cancer
Dumontet 1994 <sup>9</sup>	Case series
Aisa 2005 <sup>10</sup>	Historical control
Mori 2006 <sup>11</sup>	Historical control, oral cryotherapy 60 minutes versus 120 minutes
Sato 2006 <sup>12</sup>	No randomization, combination of cryotherapy with propantheline versus cryotherapy or propantheline alone, or no treatment.
Ohbayashi 2008 <sup>13</sup>	Not RCTs
Bhatt 2010 <sup>14</sup>	Historical control
Vokurka 2011 <sup>15</sup>	Observational study
Vokurka 2012 <sup>16</sup>	Not RCTs
Vokurka 2014 <sup>17</sup>	Not RCTs
Batlle 2014 <sup>18</sup>	Retrospective cohort study
Svanberg 2012 <sup>19</sup>	No primary and secondary outcomes

Note: RCTs: randomized controlled trials

## References

- 1 Mahood DJ, Dose AM, Loprinzi CL, Veeder MH, Athmann LM, Therneau TM *et al.* Inhibition of fluorouracil-induced stomatitis by oral cryotherapy. *J Clin Oncol*, 1991;9(3):449-452.
- 2 Rocke LK, Loprinzi CL, Lee JK, Kunselman SJ, Iverson RK, Finck G *et al.* A randomized clinical trial of two different durations of oral cryotherapy for prevention of 5-fluorouracil-related stomatitis. *Cancer*, 1993;72(7):2234-2238.
- 3 Cascinu S, Fedeli A, Fedeli SL, Catalano G. Oral cooling (cryotherapy), an effective treatment for the prevention of 5-fluorouracil-induced stomatitis. *Eur J Cancer B Oral Oncol*, 1994;30b(4):234-236.
- 4 Nikoletti S, Hyde S, Shaw T, Myers H, Kristjanson LJ. Comparison of plain ice and flavoured ice for preventing oral mucositis associated with the use of 5 fluorouracil. *J Clin Nurs*, 2005;14(6):750-753.
- 5 Karagozoglu S, Filiz Ulusoy M. Chemotherapy: the effect of oral cryotherapy on the development of mucositis. *J Clin Nurs*, 2005;14(6):754-765.
- 6 Baydar M, Dikilitas M, Sevinc A, Aydogdu I. Prevention of oral mucositis due to 5-fluorouracil treatment with oral cryotherapy. *J Natl Med Assoc*, 2005;97(8):1161-1164.
- 7 Papadeas E, Naxakis S, Riga M, Kalofonos C. Prevention of 5-fluorouracil-related stomatitis by oral cryotherapy: a randomized controlled study. *Eur J Oncol Nurs*, 2007;11(1):60-65.
- 8 Katranci N, Ovayolu N, Ovayolu O, Sevinc A. Evaluation of the effect of cryotherapy in preventing oral mucositis

- associated with chemotherapy - a randomized controlled trial. *Eur J Oncol Nurs*, 2012;16(4):339-344.
- 9 Dumontet C, Sonnet A, Bastion Y, Salles G, Espinouse D, Coiffier B. Prevention of high dose L-PAM-induced mucositis by cryotherapy. *Bone Marrow Transplant*, 1994;14(3):492-494.
- 10 Aisa Y, Mori T, Kudo M, Yashima T, Kondo S, Yokoyama A *et al*. Oral cryotherapy for the prevention of high-dose melphalan-induced stomatitis in allogeneic hematopoietic stem cell transplant recipients. *Support Care Cancer*, 2005;13(4):266-269.
- 11 Mori T, Yamazaki R, Aisa Y, Nakazato T, Kudo M, Yashima T *et al*. Brief oral cryotherapy for the prevention of high-dose melphalan-induced stomatitis in allogeneic hematopoietic stem cell transplant recipients. *Support Care Cancer*, 2006;14(4):392-395.
- 12 Sato A, Saisho-Hattori T, Koizumi Y, Minegishi M, Iinuma K, Imaizumi M. Prophylaxis of mucosal toxicity by oral propantheline and cryotherapy in children with malignancies undergoing myeloablative chemo-radiotherapy. *Tohoku J Exp Med*, 2006;210(4):315-320.
- 13 Ohbayashi Y, Imataki O, Ohnishi H, Iwasaki A, Ogawa T, Inagaki N *et al*. Multivariate analysis of factors influencing oral mucositis in allogeneic hematopoietic stem cell transplantation. *Ann Hematol*, 2008;87(10):837-845.
- 14 Bhatt V, Vendrell N, Nau K, Crumb D, Roy V. Implementation of a standardized protocol for prevention and management of oral mucositis in patients undergoing hematopoietic cell transplantation. *J Oncol Pharm Pract*, 2010;16(3):195-204.
- 15 Vokurka S, Bystricka E, Scudlova J, Mazur E, Visokaiova M, Vasilieva E *et al*. The risk factors for oral mucositis and the effect of cryotherapy in patients after the BEAM and HD-I-PAM 200 mg/m<sup>2</sup> autologous hematopoietic stem cell transplantation. *Eur J Oncol Nurs*, 2011;15(5):508-512.
- 16 Vokurka S, Svoboda T, Jungova A, Karas M, Koza V. Oral cryotherapy can significantly reduce oral mucositis but not acute GVHD incidence in Flu/Mel conditioning allo-SCT. *Bone Marrow Transplant*, 2012;47(5):739-741.
- 17 Vokurka S, Chvojkova I, Svoboda T, Brandejsova R, Jungova A, Bystricka E *et al*. The impact of oral cryotherapy and oral and gastrointestinal mucositis after autologous stem cell transplantation. *Eur J Oncol Nurs*, 2014;18(2):228-229.
- 18 Batlle M, Morgades M, Vives S, Ferra C, Oriol A, Sancho JM *et al*. Usefulness and safety of oral cryotherapy in the prevention of oral mucositis after conditioning regimens with high-dose melphalan for autologous stem cell transplantation for lymphoma and myeloma. *Eur J Haematol*, 2014.
- 19 Svanberg A, Ohrn K, Birgegard G. Five-year follow-up of survival and relapse in patients who received cryotherapy during high-dose chemotherapy for stem cell transplantation shows no safety concerns. *Eur J Cancer Care (Engl)*, 2012;21(6):822-828.