

Web references

- w1 Steinau M, Reddy D, Sumbry A, et al. Oral sampling and human papillomavirus genotyping in HIV-infected patients. *J Oral Pathol Med* 2012;41:288–91. doi:10.1111/j.1600-0714.2011.01093.x
- w2 Kreimer AR, Pierce Campbell CM, Lin H-Y, et al. Incidence and clearance of oral human papillomavirus infection in men: the HIM cohort study. *The Lancet* 2013;382:877–87. doi:10.1016/S0140-6736(13)60809-0
- w3 Bouvard V, Baan R, Straif K, et al. A review of human carcinogens--Part B: biological agents. *Lancet Oncol* 2009;10:321–2.
- w4 Videla S, Darwich L, Cañadas M-P, et al. Natural history of human papillomavirus infections involving anal, penile, and oral sites among HIV-positive men. *Sex Transm Dis* 2013;40:3–10. doi:10.1097/OLQ.0b013e31827e87bd
- w5 Parisi SG, Cruciani M, Scaggiante R, et al. Anal and oral human papillomavirus (HPV) infection in HIV-infected subjects in northern Italy: a longitudinal cohort study among men who have sex with men. *BMC Infect Dis* 2011;11:150. doi:10.1186/1471-2334-11-150
- w6 Edelstein ZR, Schwartz SM, Hawes S, et al. Rates and determinants of oral human papillomavirus infection in young men. *Sex Transm Dis* 2012;39:860–7. doi:10.1097/OLQ.0b013e318269d098
- w7 Herrero R, Quint W, Hildesheim A, et al. Reduced prevalence of oral human papillomavirus (HPV) 4 years after bivalent HPV vaccination in a randomized clinical trial in Costa Rica. *PLoS ONE* 2013;8:e68329. doi:10.1371/journal.pone.0068329