

Title page

Disrupted resting-state attentional networks in T2DM patients

Authors:

Wenqing Xia^{1,2,#}, Shaohua Wang^{3,#,*}, Hengyi Rao², Andrea M Spaeth⁴, Pin Wang¹, Yue Yang¹,
Rong Huang¹, Rongrong Cai¹, Haixia Sun¹

[#] These authors contributed equally to the preparation of this manuscript.

Author Affiliations:

¹ Medical school of Southeast University, No.87 Dingjiaqiao Road, Nanjing, China, 210009;

² Center for functional Neuroimaging, University of Pennsylvania, 3710 Hamilton Walk,
Philadelphia, PA, USA, 19104;

³ Department of Endocrinology, affiliated ZhongDa Hospital of Southeast University, No.87
Dingjiaqiao Road, Nanjing, China, 210009;

⁴ Center for Sleep and Circadian Neurobiology, Perelman School of Medicine, University of
Pennsylvania, 3710 Hamilton Walk, Philadelphia, PA, 19104

***Corresponding Author:**

Shaohua Wang, Ph.D, Prof.

Department of Endocrinology, affiliated Zhongda Hospital of Southeast University

No.87 Dingjiaqiao Road, Nanjing, China, 210009

E-mail: gyjwsh@126.com; Tel: 0086-25-83272261.

Supplementary Table 1. Therapeutic agents of the T2DM patients.

Therapeutic agents	Frequency (n)
Insulin	15
Biguanides	16
Glucosidase Inhibitors	6
Sulfonylureas	10
Non-Sulfonylurea	4
Thiazolidinediones	6
Only Diet and Exercise	1
