

Supplementary Table 2 Proportion of *Streptococcus pneumoniae* and *Haemophilus influenzae* serotypes by AOM episodes

HIV+ (N=15)		HIV- (N=182)		HEU (N=63)		Overall (N=260)		
n	%	n	%	n	%	n	%	
<i>Streptococcus pneumoniae</i> (N=53)								
<u>Vaccine types</u>								
1	1^	50.0	-	-	-	1	1.9	
3	-	-	1	2.7	1	7.1	2	3.8
4	-	-	3	8.1	1	7.1	4	7.5
5	-	-	1	2.7	-	-	1	1.9
6A	-	-	2	5.4	2	14.3	4	7.5
6B	1^	50.0	2	5.4	1	7.1	3+1^	7.5
9V	-	-	1	2.7	1	7.1	2	3.8
14	-	-	3	8.1			3	5.7
18C	-	-	1	2.7	-	-	1	1.9

21 April 2014

19A	1	50.0	4	10.8	1	7.1	6	11.3
19F	-	-	7+1^	21.6	4	28.6	11+1^	22.6
23F	-	-	2+2^	10.8	-	-	2+2^	7.5

Non-vaccine types

9N	-	-	1	2.7	-	-	1	1.9
15B	-	-	3	8.1	3	21.4	6	11.3
15C	-	-	1	2.7	-	-	1	1.9
16F	-	-	2+1^	8.1	-	-	2+1^	8.1
21	-	-	1	2.7	-	-	1	1.9

***Haemophilus influenzae* (N=80)**

F	1	14.3	-	-	-	-	1	1.3
Non-typeable	6	85.7	53	100	20	100	79	98.8
(NTHI)								

[^]Corresponds to two samples collected from one episode that tested positive for more than one *S. pneumoniae* serotype

21 April 2014

N: total number of episodes positive for each bacterium; n: number of episodes; %: percentage of episodes in a bacterial category positive for each serotype

HIV+: Children confirmed to be HIV-infected; HIV-: Children confirmed to be HIV-uninfected; HEU: children born to HIV-infected mothers and confirmed to be HIV-uninfected