

S4_Table. Estimation of genetic diversity after extra data exclusion steps. Recalculations of the genetic diversity were performed after removal of three species (*B. subterraneus*, *B. ruderatus* and *B. lapidarius*) and populations with non-amplifications and based on the same 8 microsatellite loci in each species.

<i>Species</i>	<i>Location</i>	<i>Year</i>	<i>n</i>	<i>A_R</i>		<i>H_E</i>	
				Mean	SE	Mean	SE
Widespread / stable							
<i>B. hortorum</i>	Gelderland	1918	8	5.428	0.845	0.720	0.081
	Overijssel	1918	7	5.515	0.429	0.779	0.025
	Z-Holland	1923	7	5.648	0.468	0.787	0.023
<i>B. pratorum</i>	Overijssel	1918	8	4.945	0.589	0.727	0.044
<i>B. pascuorum</i>	Limburg	1918	9	4.962	0.582	0.694	0.085
	N-Holland	1924	9	4.777	0.692	0.702	0.072
	Overijssel	1918	8	5.035	0.704	0.685	0.089
	Gelderland	1925	7	5.250	0.457	0.733	0.041
	Total		63	5.195	0.118	0.728	0.014
Restricted / declining							
<i>B. humilis</i>	Gelderland	1926	8	3.546	0.410	0.574	0.078
	Limburg	1918	8	3.182	0.363	0.522	0.072
<i>B. sylvarum</i>	Limburg	1918	6	3.821	0.610	0.601	0.089
	Limburg	1920	5	3.286	0.565	0.589	0.069
	Subtotal		27	3.459	0.143	0.572	0.017
Widespread / declining							
<i>B. muscorum</i>	Limburg	1918	7	3.603	0.640	0.516	0.109
	Overijssel	1918	8	4.360	0.517	0.613	0.078
<i>B. ruderarius</i>	Limburg	1918	7	4.149	0.594	0.620	0.102

<i>Species</i>	Location	Year	<i>n</i>	<i>A_R</i>		<i>H_E</i>			
				Mean	SE	Mean	SE		
<i>B. veteranus</i>	N-Holland	1924	5	3.750	0.697	0.610	0.089		
	Overijssel	1918	6	3.663	0.792	0.566	0.107		
	Limburg	1918	7	4.153	0.423	0.619	0.064		
	<i>Subtotal</i>		40	3.946	0.128	0.591	0.017		
<hr/>			Total	67	3.751	0.144	0.583		
							0.014		

