

## Intratumoral diversity of telomere length in individual Neuroblastoma tumors

### Supplementary Material

#### Demographic and clinical characteristics of NB patients (N=102) and type of TL (heterogeneous/homogeneous)

	TL: “heterogeneous”	TL: “homogeneous”	P
<b>Age:</b>			
≥ 18 months	13/44 (29.6%)	31/44 (70.4%)	0.73
< 18 months	19/58 (32.8%)	39/58 (67.2%)	
<b>Stage:</b>			
1	6/13 (46.2%)	7/13 (53.8%)	0.054 <sup>§</sup>
2A/2B	13/18 (72.2%)	5/18 (27.8%)	
3	16/28 (57.1%)	12/28 (42.9%)	
4	27/31 (87.1%)	4/31 (12.9%)	
4s	7/12 (58.3%)	5/12 (41.7%)	
<b>MYCN status:</b>			
Amplified	15/44 (34.1%)	29/44 (65.9%)	0.61
Not amplified	17/58 (29.3%)	41/58 (70.7%)	
<b>Ploidy:</b>			
Diploid	16/49 (32.6%)	33/49 (67.4%)	0.79
Hyperdiploid	16/53 (30.2%)	37/53 (69.8%)	
<b>Histology:</b>			
Unfavorable histology	16/50 (32.0%)	34/50 (68.0%)	0.89
Favorable histology	16/52 (30.8%)	36/52 (69.2%)	
<b>Risk Group (N=86):</b>			
High/Intermediate	18/58 (31.0%)	40/58 (69.0%)	0.92
Low	9/28 (32.1%)	19/28 (67.9%)	

P values refer to the Pearson’s Chi-square test unless otherwise specified; <sup>§</sup> Fisher’s Exact test.