

Table S1. Demographic characteristics of study participants.

Characteristic	Pre-Donation Visit		36 Month Visit		Study States (Weighted) ¹	United States ²
	Controls (N=201)	Donors (N=203)	Controls (N=173)	Donors (N=182)		
Male	32.3%	32.0%	34.1%	34.1%	40.3%	39.8%
White ethnicity	95.0%	94.6%	95.4%	94.0%	85.7%	69.7%
Age (years)	43.1±11.9	43.4±11.3	46.5±12.3	46.7±11.3		
Age category						
18-34 years	30.9%	28.1%	23.7%	17.6%	33.4%	33.2%
35-49 years	35.3%	40.9%	32.4%	39.0%	42.8%	44.6%
50-64 years	31.3%	28.6%	39.3%	39.6%	21.9%	20.9%
≥65 years	2.5%	2.5%	4.6%	3.9%	1.8%	1.3%
BMI (kg/m ²)	26.7±4.9 (200)	26.8±4.1 (201)	27.2±4.9 (173)	27.2±4.4 (181)		
BMI category						
<25 kg/m ²	41.8%	36.5%	40.5%	35.2%		
25-30 kg/m ²	35.3%	40.9%	32.4%	39.6%		
30-40 kg/m ²	20.9%	20.7%	25.4%	23.6%		
≥40 kg/m ²	1.5%	1.0%	1.7%	1.1%		
BMI missing	0.5%	1.0%	0.0%	0.6%		
Waist circ. (cm)	87.2±12.8 (181)	88.1±12.2 (174)	88.7±13.2 (162)	89.1±12.7 (173)		
Hip circ. (cm)	102.7±10.0 (170)	103.1±9.8 (163)	104.0±9.6 (162)	103.8±8.4 (171)		

Values are mean ± standard deviation and (N in parentheses) if different from complete sample.

Abbreviations: BMI, body mass index; circ., circumference.

¹Among living kidney donors in the states with participating sites (Minnesota, Ohio, Maryland, Iowa, and California), weighted by the proportion of donors enrolled in those states.

²All living donors in the US that donated 2006-2010.