

S2 Table. Basic body data for groups.

| Group | BW (g) | LKW (g) | Plasma [Na ⁺] (mEq) | Hct (%) |
|--------------|-----------|-------------|---------------------------------------|------------|
| Control | 218 ± 8 | 1.02 ± 0.08 | 139.2 ± 0.5 | 43.2 ± 1.1 |
| LPS8 | 221 ± 7 | 1.05 ± 0.09 | 140.6 ± 0.3 | 42.1 ± 1.3 |
| LPS24 | 213 ± 9 | 1.07 ± 0.10 | 140.9 ± 0.6 | 44.6 ± 2.1 |
| LPS48 | 216 ± 10 | 1.11±0.11 | 141.1 ± 0.8 | 43.9±0.9 |
| MK-801 | 220 ± 9 | 1.04 ± 0.09 | 140.5 ± 0.3 | 41.3 ± 1.6 |
| LPS8+MK-801 | 225 ± 8 | 1.06 ± 0.10 | 140.2 ± 0.4 | 42.3 ± 1.8 |
| LPS24+MK-801 | 221 ± 10 | 0.99 ± 0.08 | 139.4 ± 0.6 | 40.6 ± 1.5 |
| LPS48+MK-801 | 218 ± 7 | 1.01 ± 0.07 | 139.1 ± 0.4 | 41.2 ± 0.9 |

Eight rats in each group. BW, body weight; LKW, left kidney weight; Hct, hematocrit.