

SUPPLEMENTARY TABLE S2. PERCENT DISTRIBUTIONS OF COVARIATES BY SELF-REPORTED REPRODUCTIVE TRACT INFECTION STATUS

<i>Covariate</i>	<i>PID Yes (n=162) %</i>	<i>Chlamydia Yes (n=628) %</i>	<i>Bacterial vaginosis Yes (n=624) %</i>	<i>Trichomonas Yes (n=530) %</i>	<i>Gonorrhea Yes (n=323) %</i>	<i>Genital herpes Yes (n=143) %</i>	<i>Genital warts Yes (n=113) %</i>
Age (years)							
23–26	24	33	28	25	29	26	20
27–30	41	33	35	34	36	34	37
≥31	36	34	37	41	35	40	43
Education							
≤High school/GED	27	19	15	20	25	18	16
Some college or technical	56	57	54	58	56	47	49
≥Bachelors	17	24	31	22	19	36	35
BMI (kg/m <sup>2</sup> )							
15–24	20	18	20	14	21	15	18
25–29	20	25	25	19	25	33	24
30–34	18	20	22	20	20	21	26
≥35	43	37	33	47	35	32	33
Alcohol at age when drinking the most <sup>a</sup> (years)							
Low	21	21	18	17	20	18	18
Moderate	22	31	36	30	31	41	36
Heavy	57	48	46	53	50	41	46
Parity							
Nulliparous	30	33	36	32	30	43	40
Parous	70	68	64	68	70	57	60
Depo-Provera							
Never used	48	52	56	50	46	52	53
Ever used	52	48	44	50	54	48	47
Age at menarche (years)							
7–10	22	19	17	20	18	20	15
11–19	78	81	83	80	82	80	85
Sex partners before age 20							
≤1	14	12	17	13	12	16	17
2–5	49	48	49	50	43	51	42
≥6	38	40	34	38	45	33	42
Age at first sex (years)							
≤14	43	38	35	41	42	40	33
15–16	32	38	37	34	36	25	34
≥17	25	24	28	25	23	35	34

<sup>a</sup>Low drinkers were those who never had 10 or more drinks in a year. Heavy drinkers were those who usually drank 6 or more drinks on days when they drank or drank 4 or more drinks per sitting at least 2 to 3 times a month. Moderate drinkers were all others. BMI, body mass index; GED, general education degree; PID, pelvic inflammatory disease.