Supplementary Table S4. Adjusted Odds Ratios for Fibroids According to Self-Reported History OF NUMBER OF DIFFERENT REPRODUCTIVE TRACT INFECTIONS AND NUMBER OF DIAGNOSES FOR EACH INFECTION

	Fibroids			
Reproductive tract infections ^a	Yes (N=363) n (%) ^a	No (N=1,293) n (%)	Age-adjusted OR (95% CI)	$\frac{\textit{Multivariable-adjusted}^{\text{b}}}{\textit{OR (95\% CI)}}$
0	101 (28)	381 (30)	1 (ref)	1 (ref)
1	103 (28)	350 (27)	1.08 (0.79–1.49)	1.19 (0.86–1.64)
2	85 (23)	238 (18)	1.32 (0.94–1.85)	1.41 (0.99–2.01)
3+	74 (20)	320 (25)	0.82 (0.58–1.15)	0.91 (0.63–1.30)
Missing		4		
PID diagnoses (n)				
0	329 (91)	1,164 (90)	1 (ref)	1 (ref)
1	28 (8)	104 (8)	0.94 (0.60–1.47)	0.98 (0.62–1.55)
2+	6 (2)	24 (2)	0.75 (0.30–1.88)	0.79 (0.31–2.02)
Missing		1		
Chlamydia diagnoses (n)				
0	238 (66)	788 (61)	1 (ref)	1 (ref)
1	80 (22)	339 (26)	0.82 (0.61–1.09)	0.86 (0.64–1.16)
2+	45 (12)	164 (13)	0.92 (0.63–1.32)	1.02 (0.70–1.50)
Missing		2		
Bacterial vaginosis diagnoses (n)				
0	216 (60)	815 (63)	1 (ref)	1 (ref)
1	50 (14)	164 (13)	1.22 (0.85–1.74)	1.23 (0.85–1.78)
2+	97 (27)	313 (24)	1.10 (0.83–1.45)	1.11 (0.83–1.49)
Missing		1		
Trichomonas diagnoses (n)				
0	247 (68)	878 (68)	1 (ref)	1 (ref)
1	76 (21)	276 (21)	0.91 (0.68–1.23)	0.93 (0.68–1.27)
2+	40 (11)	138 (11)	0.88 (0.60–1.30)	0.90 (0.60–1.35)
Missing	` ′	1	` ,	,
Gonorrhea diagnoses (n)				
0	301 (83)	1,031 (78)	1 (ref)	1 (ref)
1	41 (11)	190 (15)	0.74 (0.51–1.07)	0.81 (0.56–1.19)
2+	21 (6)	71 (6)	1.01 (0.61–1.69)	1.21 (0.71–2.04)
Missing	(-)	1	32 = (312 = 2132)	(
		-		

Any "prefer not to answer" response was coded as missing.

aGenital herpes was not included because the number of times diagnosed was not applicable, and genital warts was not included because

only 9 participants were diagnosed more than once.

^bLogistic regression model adjusted for age (continuous), education, body mass index, alcohol, menarche, parity, and Depo-Provera use. CI, confidence interval; OR, odds ratio.