SUPPLEMENTARY TABLE S2. DEMOGRAPHIC CHARACTERISTICS OF 630 TERM, PREVIOUSLY HEALT	ΉY
INFANTS WITH ACUTE RESPIRATORY ILLNESS, AND THEIR MOTHERS, WHO WERE ENROLLED	
in the Tennessee Children's Respiratory Initiative	

	<i>LRTI</i> (N=455; 72%)	URI (N=175; 28%)
Baseline demographics:		
Mother's age at infant birth in years, median (IQR)	26 (22, 30)	23 (20, 29)
Insurance:		
Private	142 (31)	19 (11)
Medicaid	280 (62)	148 (85)
None	33 (7)	8 (5)
Infant's race:		
African American	82 (18)	59 (33)
White	277 (61)	52 (30)
Hispanic	53 (12)	31 (18)
Other	42 (9)	33 (19)
Gestational age in weeks, median (IQR)	39 (38, 40)	39 (39, 40)
Infant male sex	253 (56)	99 (57)
Birth weight in grams, median (IQR)	3,289 (3,005, 3,629)	3,345 (3,062, 3,657)
Breast fed	248 (55)	108 (62)
Infant in day care	105 (23)	48 (28%)
No older siblings	82 (18%)	74 (42%)
Mother currently smoking	143 (31)	46 (27)
Maternal asthma	85 (19)	37 (21)
Acute respiratory illness characteristics:		
Infant's age at time of respiratory illness in weeks, median (IQR)	11 (6, 21)	23 (9.5, 39)
RSV positive ^a	344 (76)	25 (14)
RV positive ^a	81 (18)	81 (46)
Total outpatient	64 (14)	143 (82)
Total inpatient	391 (86)	32 (18)
ICU	40 (9)	0 (0)
ICU and intubated	19 (4)	0 (0)
Respiratory severity score total, median (IQR) ^b	6.5 (4, 8.5)	1 (0, 2)
Maximum location of care	Respiratory severity score, median (IQR)	
Outpatient $(n=207)$	1 (0, 3)	
Inpatient $(n = 423)$	6.5(4, 9)	
ICU (n=40)	9 (7.5, 12)	
ICU + intubated $(n = 19)$	9.5 (9, 12)	

Unless otherwise stated, all information is presented as N (%). ^aIncludes co-infections with other viruses not included in this table. ^bRSS is a 0–12-point scale, with higher score denoting more severe disease.