

Table S1. Two-Bio-Parent Households: Multivariate Analysis of Mother- and Father-Reported Child Externalizing Behavior as a Function of Demographic Characteristics, Family History of Externalizing Disorders, and Treatment Condition.

Parameter	Mother Report						Father Report					
	BASELINE			POSTTREATMENT			BASELINE			POSTTREATMENT		
	B (SE)	t	p	B (SE)	t	p	B (SE)	t	p	B (SE)	t	p
Intercept	62.41 (3.32)	18.81	<.0001	61.09 (3.83)	15.95	<.0001	57.07 (3.79)	15.05	<.0001	57.30 (4.16)	13.77	<.0001
Treatment/control (ref=control)												
Treatment	-1.27 (1.24)	-1.02	0.31	-5.71 (1.43)	-3.99	<.0001	-1.35 (1.31)	-1.03	0.31	-5.77 (1.44)	-4.01	<.0001
Combined hx of externalizing disorders (ref=NO first- or second-degree risk)												
First- AND second-degree risk	5.25 (1.53)	3.43	<.001	5.24 (1.77)	2.96	<.01	4.02 (1.63)	2.47	<.05	3.90 (1.78)	2.18	<.05
First-degree risk only	2.60 (2.22)	1.17	ns	2.06 (2.56)	0.81	ns	2.01 (2.44)	0.82	ns	2.14 (2.68)	0.80	ns
Second-degree risk only	1.91 (1.36)	1.40	ns	1.16 (1.57)	0.74	ns	-0.07 (1.43)	-0.05	ns	0.82 (1.56)	0.53	ns
Child's sex (ref=Female)												
Male	0.82 (1.36)	0.60	ns	0.51 (1.57)	0.32	ns	-0.97 (1.44)	-0.67	ns	2.04 (1.58)	1.29	ns
Maternal education (ref= high school or less)												
4-year college or more	0.19 (1.53)	0.13	ns	-0.12 (1.77)	-0.07	ns	1.21 (1.66)	0.73	ns	-1.24 (1.83)	-0.68	ns
Some College	-0.67 (1.63)	-0.41	ns	-0.81 (1.88)	-0.43	ns	2.82 (1.77)	1.59	ns	-0.44 (1.95)	-0.23	ns
Maternal race (ref=non-white)												
White	1.15 (2.54)	0.45	ns	-0.15 (2.93)	-0.05	ns	5.63 (2.83)	1.99	<.05	1.81 (3.10)	0.58	ns
Maternal/paternal depression at baseline^a												
BDI	0.18 (0.10)	1.78	ns	0.20 (0.12)	1.71	ns	0.28 (0.13)	2.18	<.05	0.39 (0.14)	2.71	<.01
Model Statistics	$r^2=0.085$			$r^2=0.122$			$r^2=0.104$			$r^2=0.158$		

Note: BDI = Beck Depression Inventory; hx = history; ref = reference.

^aDepression score corresponds with the reporter of child behavior in the model.

Table S2. Single-Bio-Mother Households: Multivariate Analysis of Mother-Reported Child Externalizing Behavior as a Function of Demographic Characteristics, Family History of Externalizing Disorders, and Treatment Condition

Parameter	Mother Report			POSTTREATMENT		
	B (SE)	t	p	B (SE)	t	p
Intercept	66.92 (4.90)	13.65	<.0001	67.43 (5.49)	12.29	<.0001
Treatment/control (ref=control)						
Treatment	-2.35 (2.24)	-1.05	ns	-7.94 (2.50)	-3.17	<.01
Combined hx of externalizing disorders (ref= NO first- or second-degree risk)						
First- AND second-degree risk	3.47 (2.30)	1.51	ns	0.64 (2.58)	0.25	ns
First-degree risk only	4.39 (3.57)	1.23	ns	0.38 (3.99)	0.09	ns
Second-degree risk only	2.44 (2.45)	1.00	ns	3.77 (2.74)	1.37	ns
Child's sex (ref=Female)						
Male	1.19 (2.16)	0.55	ns	-2.54 (2.42)	-1.05	ns
Maternal education (ref= high school or less)						
4-year college or more	-4.09 (2.58)	-1.58	ns	-3.93 (2.89)	-1.36	ns
Some College	-0.57 (2.32)	-0.25	ns	-3.04 (2.60)	-1.17	ns
Maternal race (ref=non-white)						
White	2.91 (3.20)	0.91	ns	4.35 (3.58)	1.22	ns
Caregiver depression at baseline						
BDI	0.08 (0.12)	0.69	ns	0.03 (0.13)	0.24	ns
Model Statistics			r ² =.155			r ² =.227

Note: BDI = Beck's Depression Inventory; hx = history; ref = reference.