

Table S5. AUC for suicide by cut-off score of Beck Depression Inventory in Korean middle-aged men.

BDI score cut-off	AUC^a (95% CI)	BDI score cut-off	AUC^a (95% CI)
5 or above	0.510(0.486-0.535)	31 or above	0.694(0.620-0.767)
6 or above	0.517(0.492-0.541)	32 or above	0.686(0.611-0.762)
7 or above	0.523(0.498-0.548)	33 or above	0.652(0.574-0.729)
8 or above	0.531(0.506-0.557)	34 or above	0.656(0.578-0.733)
9 or above	0.541(0.516-0.566)	35 or above	0.663(0.585-0.740)
10 or above	0.551(0.526-0.577)	36 or above	0.657(0.580-0.735)
11 or above	0.550(0.515-0.584)	37 or above	0.663(0.586-0.741)
12 or above	0.550(0.509-0.592)	38 or above	0.644(0.567-0.720)
13 or above	0.563(0.521-0.605)	39 or above	0.648(0.572-0.724)
14 or above	0.576(0.533-0.619)	40 or above	0.623(0.549-0.697)
15 or above	0.577(0.528-0.625)	41 or above	0.619(0.546-0.692)
16 or above	0.567(0.509-0.624)	42 or above	0.617(0.545-0.689)
17 or above	0.580(0.522-0.638)	43 or above	0.603(0.534-0.673)
18 or above	0.592(0.533-0.651)	44 or above	0.578(0.514-0.643)
19 or above	0.604(0.544-0.664)	45 or above	0.577(0.514-0.641)
20 or above	0.617(0.557-0.678)	46 or above	0.576(0.513-0.638)
21 or above	0.617(0.553-0.682)	47 or above	0.562(0.504-0.620)
22 or above	0.629(0.564-0.694)	48 or above	0.558(0.502-0.614)
23 or above	0.641(0.575-0.707)	49 or above	0.539(0.490-0.587)
24 or above	0.653(0.587-0.720)	50 or above	0.532(0.488-0.576)
25 or above	0.667(0.600-0.734)	51 or above	0.531(0.488-0.573)
26 or above	0.678(0.611-0.746)	52 or above	0.530(0.488-0.571)
27 or above	0.664(0.592-0.736)	53 or above	0.516(0.484-0.549)
28 or above	0.675(0.603-0.747)	54 or above	0.511(0.483-0.538)
29 or above	0.682(0.610-0.755)	55 or above	0.509(0.484-0.533)
30 or above	0.686(0.612-0.759)	56 or above	0.509(0.485-0.533)

AUC, area under the receiver operating characteristics curve; BDI, Beck Depression Inventory; CI, confidence interval.

a. A diagnostic test with an AUC value of 1.0 or 0.5 represents a perfect test or an uninformative test, respectively