

81 **Table S1. Strains used in this study**

Strain	Characteristics	Reference
LVS	Live vaccine strain	USAMRIID ^a
<i>ΔigIA</i>	LVS, <i>igIA</i> null mutant	(1)
<i>ΔigIC</i>	LVS, <i>igIC</i> null mutant	(2)
<i>ΔigIG</i>	LVS, <i>igIG</i> null mutant	(3)
<i>ΔigII</i>	LVS, <i>igII</i> null mutant	(3)
<i>ΔpdpE</i>	LVS, <i>pdpE</i> null mutant	(3)
<i>L. monocytogenes</i> EGD	EGD wild type strain	(4)

82 ^a USAMRIID, U.S. Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick, MD.

83

84 **References**

- 85 1. Bröms JE, Lavander M, Sjöstedt A. 2009. A conserved a-helix essential for a type VI
86 secretion-like system of *Francisella tularensis*. J Bacteriol **191**:2431-2446.
- 87 2. Lindgren H, Golovliov I, Baranov V, Ernst RK, Telepnev M, Sjöstedt A. 2004. Factors affecting
88 the escape of *Francisella tularensis* from the phagolysosome. J Med Microbiol **53**:953-958.
- 89 3. Bröms JE, Lavander M, Meyer L, Sjöstedt A. 2011. IgIG and IgII of the *Francisella*
90 pathogenicity island are important virulence determinants of *Francisella tularensis* LVS.
91 Infect immun **79**:3683-3696.
- 92 4. Mandin P, Fsihi H, Dussurget O, Vergassola M, Milohanic E, Toledo-Arana A, Lasa I,
93 Johansson J, Cossart P. 2005. VirR, a response regulator critical for *Listeria monocytogenes*
94 virulence. Mol Microbiol **57**:1367-1380.

95

96