
## 5. ADVICE FOR ALL CHILDREN TREATED AT HOME

### Give more fluids and continue feeding

- Advise caregiver to give plenty of homemade fluids such as:
  - · Clean water
  - Soup
  - Yoghurt drinks
  - Light porridge
  - ORS



#### Go to the health facility if...

- Advise to go to the health facility if the child:
  - · Cannot drink or feed
  - Has blood in stool
  - · Becomes sicker
  - · Develops any other danger sign
  - · Has not improved in 2 days




#### Sleep under a net

Advise caregiver to be sure the child sleeps under a mosquito net.



#### Follow up child in 3 days

Decide if you (the VHT member) will go to the child or if the caregiver will bring the child to you.





# 6. ROUTINE CARE FOR THE NEWBORN

#### Keeping the baby warm

Advise the caregiver on the following:

- Wrap the baby in warm, dry clothes, including the head and feet.
- Put the baby in skin-to-skin contact with the mother, as shown in the picture.
- Delay the baby's first bath until after 24 hours.





#### Breastfeeding exclusively

Advise the mother on the following:

- Start breastfeeding immediately after birth.
- Feed the baby on only breast milk, on demand, at least 8 times a day.
- Ensure the baby is well positioned and attached on the mother's breast.

#### Skin and cord care

Advise the caregiver on the following:

Wash hands before handling the baby.



Do not apply anything on the cord. Leave the cord dry and



Bathe the baby with clean soap and water.











