

Supplemental Data


Acetylbritannilactone Modulates MicroRNA-155-Mediated Inflammatory Response in Ischemic Cerebral Tissues

Ya Wen,^{1,2,3} Xiangjian Zhang,^{1,2,3} Lipeng Dong,^{1,2} Jingru Zhao,^{1,2} Cong Zhang,^{1,2} and Chunhua Zhu^{1,2,3}

¹Department of Neurology, Second Hospital of Hebei Medical University, Shijiazhuang, Hebei, PR China; ²Hebei Key Laboratory for Neurology, Shijiazhuang, Hebei, PR China; and ³Hebei Collaborative Innovation Center for Cardio-cerebrovascular Disease, Shijiazhuang, Hebei, PR China

Online address: <http://www.molmed.org>

The Feinstein Institute for Medical Research North Shore LIJ
Empowering Imagination. Pioneering Discovery.®



Supplementary Figure S1. (A) The widespread of miR-155 in the brain after the stereotactic injections of pAd-miR-155 into LV. Left, miR-155 was widely expression in the whole mouse brain, especially in cortex and hippocampus, as assessed by the fluorescence intensity of the transfected pAd-miR-155 tagged with GFP. Right, bright field of the brain after the stereotactic injections of pAd-miR-155. (B) miR-155 levels were increased in the brain after 24 h of delivery of pAd-miR-155. Western blotting means the expression of the transfected pAd-miR-155 tagged with GFP using anti-GFP antibody, showing that the introduced pAd-miR-155 tagged with GFP was highly expressed in the brain tissues after 24 h of delivery of pAd-miR-155.