

SUPPLEMENTARY TABLE S1. RESULTS OF THE TWO-WAY ANOVA TESTS REFERRED TO IN THE TEXT

GFP area	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	26	5	5.4709	0.0025
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Side	2	0.8106	0.4587
	Time	2	10.2463	0.0009
	Side	1	4.6499	0.0434
Collagen scar area	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	45	9	1.3438	0.2508
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Group	4	0.0907	0.9848
	Time	4	2.7147	0.0455
	Group	1	0.5085	0.4805
Fibrinoid scar area	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	45	9	3.7983	0.002
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Group	4	1.5362	0.2131
	Time	4	7.358	0.0002
	Group	1	0.2069	0.652
Collagen scar distance	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	45	9	2.8166	0.0134
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Group	4	0.8315	0.5143
	Time	4	5.1995	0.0021
	Group	1	0.001	0.9744
Fibrinoid scar distance	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	45	9	1.4093	0.2218
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Group	4	0.286	0.8851
	Time	4	2.7038	0.0461
	Group	1	0.1248	0.726
GFAP astrocytes distance	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	45	9	2.9026	0.0113
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Group	4	0.9525	0.4456
	Time	4	5.3581	0.0018
	Group	1	3.7855	0.0598
NF axons distance	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	45	9	1.1003	0.3879
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Group	4	1.0059	0.4176
	Time	4	0.8063	0.5296
	Group	1	2.9366	0.0954
CSPG perineuronal nets distance	Total <i>N</i>	Model DF	F Ratio	<i>p</i> -value
	45	9	0.668	0.7316
	Effect variable	DF	F Ratio	<i>p</i> -value
	Time × Group	4	0.2449	0.9108
	Time	4	1.1596	0.3452
	Group	1	0.3323	0.568

CSPG, chondroitin sulfate proteoglycan; GFAP, glial fibrillary acidic protein; GFP, green fluorescent protein; NF, neurofilament.