Drug Safety

Actual Use of Medications Prescribed During Pregnancy: A Cross-Sectional Study using data from a Population Based Congenital Anomaly Registry

L de Jonge, HEK de Walle, LTW de Jong-van den Berg, IM van Langen, MK Bakker

Electronic Supplementary Material #2: Compliance rates calculated according to the standard definition, in a selected population including only live births and still births and according to the strict compliance definition

	"standard" ~ definition of	Compliance rate among LB and SB population,	Strict* 'compliance'
	compliance	excluding SA and TOPFAs	definition
F:	compliance	excluding 3A and TOFTAS	definition
First trimester			
Total number of women	129	107	129
medications for chronic use	0.84 (0.74-0.95)^	0,83 (0,71-0,96)^	0,82 (0,71-0,93)^
medications for short-term and occasional use	0.86 (0.78-0.94)^	0,85 (0,76-0,94)^	0,84 (0,76-0,93)^
medications for pregnancy related symptoms	0.92 (0.82-0.96)#	0,92 (0,81-0,97)#	0,88 (0,80-0,96)^
Entire pregnancy			
Total number of women	202	172	202
medications for chronic use	0.92 (0.86-0.98)^	0,92 (0,84-0,97)#	0,91 (0,84-0,97)^
medications for short-term and occasional use	0.90 (0.85-0.95)^	0,89 (0,83-0,95) ^	0,89 (0,84-0,94)^
medications for pregnancy related symptoms	0.95 (0.90-0.99)^	0,96 (0,90-0,98) #	0,92 (0,87-0,97)^

[~] standard definition of compliance: If a mother confirmed that she had taken **at least one** of the prescriptions of a specific medication. If a medication was prescribed more than once and the mother had taken just one prescription, she was still counted as a compliant user for that medication.

^{*} strict definition of compliance: if a mother confirmed that she had taken **all** of the prescriptions of a specific medication. If a medication was prescribed more than once but not all the prescriptions had been taken, she was not counted as a 'compliant user'.

[^] Wald-method was applied

[#] Wilson-method was applied