

Time-lapse Raman imaging of osteoblast differentiation

Supplementary Information

Aya Hashimoto^a, Yoshinori Yamaguchi^{a,b,*}, Liang-da Chiu^a, Chiaki Morimoto^c,

Katsumasa Fujita^a, Masahide Takedachi^c, , Satoshi Kawata^a,

Shinya Murakami^c, and Eiichi Tamiya^{a,*}

^a Department of Applied Physics, Graduate School of Engineering, Osaka University,
2-1 Yamadaoka, Suita, Osaka, 565-0871, Japan

^b Department of Physics, Graduate School of Science, East China University of
Science and Technology, 130 Meilong Rd., Shanghai, 200237, China

^c Department of Periodontology, Graduate School of Dentistry, Osaka University,
1-8 Yamadaoka, Suita, Osaka, 565-0871, Japan

*Corresponding authors:

Yoshinori Yamaguchi, yoshi.yamaguchi@ap.eng.osaka-u.ac.jp

Eiichi Tamiya, tamiya@ap.eng.osaka-u.ac.jp

Figure Legends

Figure S1. Noise reduction of Raman spectra by SVD calculation. (A) Raw and processed data of Raman spectra obtained from KUSA-A1. (B) Plots of the first ten basis vectors resulting from SVD calculation. The blue-letter values indicate contribution factor of the basis vectors.