

Table 4. Correlations of NEI-VFQ-25 Scores with Symptoms and Clinical Parameters in patients with Ocular Graft- versus-host Disease after Adjustment for Age and Sex

NEI-VFQ-25 Scales	OSDI	SANDE	CFS	TBUT	Schirmer	BCVA
Composite score	-0.8*	-0.5*	-0.3*	0.2	-0.1	-0.5*
Ocular pain	-0.7*	-0.6*	-0.2	0.2	0.1	-0.3*
VS Role difficulties	-0.7*	-0.4*	-0.2	0.2	-0.0	-0.3*
VS Mental function	-0.8*	-0.5*	-0.4*	0.2	0.1	-0.4*
General vision	-0.5*	-0.2*	-0.4*	0.2	0.1	-0.4*
Near activities	-0.8*	-0.5*	-0.4*	0.1	0.1	-0.4*
Distance activities	-0.7*	-0.4*	-0.4*	0.2	-0.1	-0.5*
Driving	-0.5*	-0.2*	-0.2	0.1	0.0	-0.4*
VS Dependency	-0.5*	-0.4*	-0.3*	0.1	-0.2	-0.5*
Peripheral vision	-0.4*	-0.2	-0.2	0.1	-0.3*	-0.2*
VS Social function	-0.5*	-0.4*	-0.1	0.1	-0.2	-0.5*
Color vision	-0.3*	-0.2	-0.1	-0.1	-0.3*	-0.1
General Health	-0.2	-0.2	-0.1	0.1	-0.1	-0.2

NEI-VQ-25: 25-item National Eye Institute Visual Function Questionnaire; SANDE: Symptoms Assessment in Dry Eye; OSDI: Ocular Surface Disease Index; CFS: corneal fluorescein staining; TBUT: tear break-up time; BCVA: best-corrected visual acuity; VS: vision specific. Higher scores in the NEI-VFQ-25 indicate better quality of life (note the negative correlations with OSDI, SANDE, CFS, and BCVA). *P<0.05