	PI Completers (total $n = 150$)	Non-Completers (total <i>n</i> = 71)	P
Degree (%)	(n = 150)	(n = 71)	
DO	9	9	
MD	69	70	0.020
NP	14	16	0.928
PA	8	6	
Specialty (%)	(n = 150)	(n = 71)	
Diabetes	3	6	
Endocrinology	7	4	
Family practice	33	35	0.0214
General practice	3	4	0.831†
Internal medicine	45	42	
Other*	9	8	
Primary practice type (%)	(n = 150)	(n = 71)	
Solo practice	27	35	
Group practice < 5	27	25	
Group practice > 5	21	17	
Institutionally salaried (hospital/clinic)	21	16	0.535†
Institutionally salaried (health maintenance organization/ managed care/insurance company)	1	0	
Other (please specify)	4	7	
Years in practice (%)	(n = 150)	(n = 71)	
<1	1	0	
1–10	39	28	1
11–20	32	41	0.450†
21–30	19	21	
> 30	9	10	
Patients with diabetes seen each week	(n = 150)	(n = 71)	
Mean number of patients	23	27	0.210
Having a dedicated individual for diabetes education within the office (%)	(n = 150)	(n = 71)	
Yes	47	39	0.270

continued on p. 2

1

	PI Completers (total $n = 150$)	Non-Completers (total $n = 71$)	P
Frequency of discussion of diet and exercise plans with measurable goals (%)	(n = 150)	(n = 71)	
Never	1	0	
Sometimes	13	14	0.5504
Frequently	47	38	0.558†
Always	40	48	
Frequency of documentation of discussions about diet and exercise in the patient chart (%)	(n = 149)	(n = 71)	
Never	1	0	
Sometimes	28	32	0.4621
Frequently	42	32	0.463†
Always	30	35	1
Frequency of referral of diabetes patients to a certified diabetes educator or recognized disease-state-management education program (%)	(n = 150)	(n = 71)	
Never	3	9	
Sometimes	48	35	0.179†
Frequently	35	41	
Always	14	16	
Frequency of documentation of these referrals in the patient chart (%)	(n = 145)	(n = 65)	
Never	3	0	
Sometimes	19	9	0.109†
Frequently	20	29	
Always	59	62	
Frequency of specific action taken when your diabetes patients are not at blood pressure or lipid goals (%)	(n = 150)	(n = 71)	
Never	0	0	0.376†
Sometimes	6	3	
Frequently	43	38	
Always	51	59	

continued on p. 3

	PI Completers (total <i>n</i> = 150)	Non-Completers (total <i>n</i> = 71)	P
Frequency of documentation of these actions in the patient chart (%)	(n = 150)	(n = 71)	
Never	0	0	
Sometimes	6	3	0.336†
Frequently	31	25	
Always	63	72	
Percentage of patients with type 2 diabetes who receive a comprehensive foot exam, including neurologic and vascular testing, at least once every 12 months (%)	(n = 150)	(n = 71)	
< 10%	5	9	
10-25%	8	7	
26-50%	11	17	0.5224
51–75%	19	20	0.532†
76–90%	21	13	
>90%	37	35	
Frequency of foot exam results documented in the medical chart (%)	(n = 150)	(n = 71)	
Never	1	6	0.226†
Sometimes	17	16	
Frequently	29	34	
Always	53	45	
Percentage of your type 2 diabetes patients at goal who have A1C values measured at least once every 6 months (%)	(n = 150)	(n = 71)	
< 10%	1	0	0.912†
10-25%	2	1	
26-50%	1	3	
51–75%	11	13	
76–90%	21	24	
>90%	63	59	

continued on p. 4

Supplemental Table 1S. Self-Assessment Responses by PI Completers and Non-Completers, continued				
	PI Completers (total <i>n</i> = 150)	Non-Completers (total <i>n</i> = 71)	P	
Frequency of specific action taken when diabetes patients are not meeting their A1C goals (%)	(n = 150)	(n = 71)		
Never	0	0		
Sometimes	4	4	0.967†	
Frequently	43	41		
Always	53	55		
Frequency of documentation of action taken when diabetes patients are not meeting their A1C goals (%)	(n = 150)	(n = 71)		
Never	0	0		
Sometimes	3	1	0.634†	
Frequently	19	23		
Always	77	76		

^{*}Includes all specialties for which there was ≤ 1 participant among Partial and Non-Completers or PI Completers. Other specialties include Allergy, Anesthesiology, Cardiovascular Disease, Emergency Medicine, Family Practice/Sports Medicine, Gynecology, IM—Diagnostic Lab Immunology, Infectious Diseases, Internal Medicine—Geriatrics, Internal Medicine—Pediatrics, Neurology, Obstetrics & Gynecology, Other Specialty, Psychiatry, Pulmonary Disease, Sleep Medicine, and Unspecified Specialty.

[†]More than 10% of cells had expected frequencies < 5.