

Performance quality scoring

Task	2 points	1 point	0 points
Safe approach	Looked around for any danger	Done too flighty	Not done or not ratable
Assess responsiveness	Called loud <i>and</i> stimulated	Only called <i>or</i> stimulated	Not done or very poor
Open airway	Infant: head in neutral position, chin lift Toddler: head tilt, chin lift	Partially done incorrectly	Not done or very poor
Assess breathing	Look, listen and feel (own cheek over patients' mouth, hand on patients' upper abdomen or chest, looks downward to chest)	Does not assess <i>all three</i> dimensions	Not done or very poor
Initial rescue breaths	Five sufficient rescue breaths with visible chest rise and correct head position	Partially done insufficiently	Not done or very poor
Assess circulation	Complete inspection of signs of life (movements) and feels carotid pulse	Feels other pulse than carotid, does not look for movements	Not done or very poor
CPR	CV ratio - 15 chest compressions in alteration with two rescue breaths	Partially done incorrectly	Not done or very poor
	Chest compression procedure - Pressure point lower half of sternum - Compression at least one-third of the depth of the chest - Complete release after each compression	Partially done incorrectly	Not done or very poor
	Chest compression technique: Infant: two or three fingers Toddler: one or two hands	Partially done incorrectly	Not done or very poor
	Rescue breaths Sufficient with visible chest rise and correct head position	Partially done insufficiently	Not done or very poor
	Efficient transitions between chest compressions and rescue breaths, e.g. positioning, fixed head position during chest compressions	Few interruptions during CPR	Many interruptions during CPR