

Growth differentiation factor-15: a p53- and demethylation-upregulating gene represses cell proliferation, invasion, and tumorigenesis in bladder carcinoma cells

Ke-Hung Tsui^{1,3,#}, Shu-Yuan Hsu^{2,#}, Li-Chuan Chung², Yu-Hsiang Lin¹, Tsui-Hsia Feng⁴, Tzu-Yi Lee², Phei-Lang Chang^{1,3}, & Horng-Heng Juang^{2,*}

¹Department of Urology, Chang Gung Memorial Hospital, Kwei-Shan, Tao-Yuan, Taiwan, ROC

²Department of Anatomy, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan, ROC

³Department of Traditional Chinese Medicine, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan, ROC

⁴School of Nursing, College of Medicine, Chang Gung University, Kwei-Shan, Tao-Yuan, Taiwan, ROC

Supplemental Figure 1




Figure Supplement 1. Modulation of GDF15 on p53 expression in human bladder carcinoma RT4 cells. Expressions of p53, GDF15, and β -actin in mock-knockdown RT4 (RT4-COLsi) and GDF15 knockdown RT4 (RT4-GDF15si) cells were determined by immunoblotting assays.