Table S4: Fold change for analysis of Notch activation by co-culture and immobilized ligand.

(A) Comparison of the number of genes upregulated after ligand-activation and downregulated with DAPT treatment, using noted cutoffs, in both co-culture and immobilized ligand assays, (B) gene list of (A) with comparison groups as in Figure 3E and red text indicating repeated gene listing, and (C) comparison of the number of genes up- and downregulated, using noted cutoffs, in MDA-MB-231 cultured with 3T3GFP or Fc.

Α

Comparison: Co-culture vs. Im	mobilized Ligand	Gene Expression	FC>2 # of Genes
MDA231/3T3DLL4 vs. MDA231/3T3GFP (-DAPT)	vs. MDA231/Fc-DLL4 vs. MDA231/Fc (-DAPT)	Upregulated	3
"Notch Signature" in co-culture	vs. "Notch Signature" on immobilized ligand	In common	1
MDA231/3T3DLL4 (+DAPT) vs. MDA231/3T3DLL4 (-DAPT)	vs. MDA231/Fc-DLL4 (+DAPT) vs. MDA231/Fc-DLL4 (-DAPT)	Downregulated	10

В

3 common (2 unique) genes upregulated by DLL4 (101 vs. 47 genes)

FAM45B family with sequence similarity 45, member B (pseudogene)

FAM45B family with sequence similarity 45, member B (pseudogene)

DYRK4 dual-specificity tyrosine-(Y)-phosphorylation regulated kinase 4

1 common "Notch Signature" gene (63 vs. 29 genes)

SCARNA4 small Cajal body-specific RNA 4

10 common genes downregulated by DAPT (101 vs. 55 genes)

TRIM34 tripartite motif containing 34 SMG1P5 SMG1 pseudogene 5

RNASEK-C17orf49 RNASEK-C17orf49 readthrough

LOC101928837 uncharacterized ncRNA LOC101928571 ZNF649 antisense RNA 1 INSL4 insulin-like 4 (placenta)

HES4 hes family bHLH transcription factor 4
FRMD8P1 FERM domain containing 8 pseudogene 1

ECSCR endothelial cell surface expressed chemotaxis and apoptosis regulator

COX7B2 cytochrome c oxidase subunit VIIb2

C

Comparison: GFP vs. Fc		Gene Expression	FC>2
			# of Genes
MDA231/3T3GFP	vs. MDA231/Fc	Upregulated	352
		Downregulated	97