

Supplemental Data for:

**Unique Toll-Like Receptor 4 Activation by NAMPT/PBEF Induces
NFκB Signaling and Inflammatory Lung Injury**

Sara M. Camp^{*1}, Ermelinda Ceco^{*2}, Carrie L. Evenoski^{*1}, Sergei M. Danilov³, Tong Zhou¹, Eddie T. Chiang¹, Liliana Moreno-Vinasco², Brandon Mapes², Jieling Zhao⁴, Gamze Gursoy⁴, Mary E. Brown², Djanybek M. Adyshev², Shahid S. Siddiqui², Hector Quijada¹, Saad Sammani², Eleftheria Letsiou², Laleh Saadat², Mohammed Yousef⁵, Ting Wang¹, Jie Liang⁴, Joe G. N. Garcia^{§1}

¹Department of Medicine and Arizona Respiratory Center, The University of Arizona;

²Institute for Personalized Respiratory Medicine, University of Illinois at Chicago;

³Department of Anesthesiology, University of Illinois at Chicago

⁴Department of Bioengineering, University of Illinois at Chicago

⁵Life Science Group, Bio-Rad Laboratories, Inc.

*Contributed equally to this work

§Corresponding Author:

Joe G. N. Garcia, MD

Senior Vice President for Health Sciences

Arizona Health Sciences Center

University of Arizona

1295 N. Martin Ave., Tucson AZ 85721-0202

Phone: (520) 626-1197

Fax: (520) 626-1460

Email: skipgarcia@email.arizona.edu

Running Title: Toll-Like Receptor 4 Activation by NAMPT/PBEF

Supplement Figure 1. Isotype control immunohistochemistry staining of p-NFκB in lung tissues. Either SB (control, top) or VILI (bottom) mouse lung sections were stained with the same immunohistochemistry protocol/reagents as previously reported minus the selective p-NFκB primary antibody, which was replaced with a non-selective IgG control antibody. No clear positive staining were identified in any of these slides. Representative images shown. Scale bar = 100μM.

IHC Isotype Controls

