

SUPPLEMENTARY DATA

Circulating TGF- β 1-regulated miRNAs and the risk of rapid progression to ESRD in type 1 diabetes

Marcus G. Pezzolesi PhD, MPH^{1,2}, Eiichiro Satake MD, PhD^{1,2}, Kevin P. McDonnell BS¹, Melissa Major BS¹, Adam M. Smiles MS¹, and Andrzej S. Krolewski MD, PhD^{1,2}

Supplementary Table 1A. Summary of TGF- β 1-regulated miRNAs Detected ($Cq \leq 30.0$) in Plasma from Normoalbuminuric Controls that were Not Selected for Examination in Rapid Progressors and Non-Progressors.

miRNA	Cq
let-7b-3p	28.66
miR-29b-3p	26.19
miR-29b-1-5p	29.52
miR-200a-5p	29.44
miR-200c-3p	29.95
miR-377-3p	29.33
miR-382-5p	27.20
miR-192-5p	27.37
miR-491-5p	29.78

SUPPLEMENTARY DATA

Supplementary Table 1B. Summary of TGF- β 1-regulated miRNAs Not Highly Detectable ($C_q > 30.0$) in Plasma from Normoalbuminuric T1D Control Patients.

miRNA	C_q
miR-21-3p	30.13
miR-29a-5p	32.63
miR-29b-2-5p	33.96
miR-29c-5p	30.31
miR-141-3p	31.84
miR-141-5p	33.88
miR-192-3p	31.21
miR-200a-3p	32.35
miR-200b-3p	31.09
miR-200b-5p	30.46
miR-200c-5p	32.70
miR-215-5p	32.05
miR-216a-5p	33.75
miR-216b-5p	33.25
miR-217	31.78
miR-296-5p	30.97
miR-298	33.23
miR-377-5p	31.42
miR-382-3p	>35.00