

SUPPLEMENTARY TABLE S1. MULTIVARIATE LOGISTIC MODELS FOR COMPLIANCE WITH SCREENING RECOMMENDATIONS IN U.S. ADULT WOMEN BY CARDIOVASCULAR DISEASE STATUS

<i>Disease/condition</i>	<i>Odds ratio (95% confidence interval) of disease vs. without disease</i>			
	<i>Pap smear</i>		<i>Mammogram</i>	
	<i>Model 1^a</i>	<i>Model 2^b</i>	<i>Model 1^a</i>	<i>Model 2^b</i>
Hypertension	1.08 (0.91–1.29)	0.97 (0.81–1.16)	1.24 (1.10–1.40)	1.18 (1.05–1.33)
Myocardial infarction	0.37 (0.22–0.64)	0.27 (0.16–0.47)	0.99 (0.72–1.36)	0.96 (0.71–1.29)
Stroke	0.68 (0.42–1.10)	0.62 (0.38–1.04)	1.28 (0.97–1.69)	1.12 (0.84–1.50)

Data Source: Disease Control and Prevention, National Center for Health Statistics (CDC/NCHS), National Health Interview Survey, 2013.

For the analyses on individual cardiovascular disease (CVD) type, women with other types of CVD were excluded from the comparison group.

^aVariables in Model 1 included disease status, age group, race, immigration status, region of residence, marital status, education level, and family income to poverty threshold ratio.

^bVariables in Model 2 further included variables from Model 1 and health insurance coverage, usual place of care, and visiting an obstetrician/gynecologist in the past year.