


Supplementary Figure 1: Generation of *cGAS*-deficient mice

(A) Schematic of the wild-type and targeted *cGAS* (*Mb21d1*) locus. Exons are indicated with solid black squares and primers used for genotyping PCR are indicated with arrows. The targeted "knockout first" locus contains a strong splice acceptor (SA) in the LacZ cassette, resulting in a truncated *cGAS* transcript in which exon 1 is spliced into the LacZ cassette. The majority of *cGAS*^{-/-} mice used in this manuscript carry this "knockout first" null allele of *cGAS*, including all mice crossed to *Trex1*^{-/-} mice. The *cGAS*^{-/-}*Trex1*^{+/+} mice used as controls in Figures 2 and 3 carry the final null *cGAS* allele that was generated as follows: mice with a conditional *cGAS*^{FLOX} allele were generated by crossing to FLPeR mice (1) to remove the FRT-flanked LacZ and Neo cassettes; mice with the final null *cGAS* allele were generated by crossing *cGAS*^{FLOX} mice to Mox2-Cre mice (2) for germline excision of the LoxP-flanked exon 2. The *cGAS*^{FLOX} and final *cGAS* alleles were detected with the following primers: FLOX Fwd, AAGGCGCATAACGATAACCACG; FLOX Rev, GCTGGGTCTAGATATCTCGAC; and KO Rev, AAGGGCACTGAGCCTCTGAG. (B) Genotyping PCR to distinguish *cGAS*^{+/+}, *cGAS*^{+/-}, and *cGAS*^{-/-} mice carrying the "knockout first" *cGAS* allele.



Supplementary Figure 2: Characterization of the IFN response in cGAS-deficient cells

(A) Quantification of *Ifnb* mRNA induction in bone marrow-derived macrophages from *cGAS*^{+/+}, *cGAS*^{+/-}, *cGAS*^{-/-}, or *Tmem173*^{-/-} (STING-deficient) mice transfected with calf-thymus DNA (CT-DNA), ISD 100-mer oligonucleotides (3), or RIG-I ligand (4) for 2h, 4h, or 8h as indicated. **(B)** Quantification of type I IFN production (right panel) using an IFN bioassay with supernatants from bone marrow-derived macrophages stimulated with the indicated ligands as in (A). **(C)** Quantification of *Ifnb* mRNA induction in mouse embryonic fibroblasts (MEFs) from *cGAS*^{+/+}, *cGAS*^{-/-}, or *Tmem173*^{-/-} (STING-deficient) embryos stimulated as in (A). **(D)** Quantification of type I IFN production (right panel) using an IFN bioassay with supernatants from MEFs stimulated with the indicated ligands as in (C). Data are representative of two experiments.

Supplementary References

1. Farley, F. W., P. Soriano, L. S. Steffen, and S. M. Dymecki. 2000. Widespread recombinase expression using FLP_{er} (flipper) mice. *Genesis* 28: 106-110.
2. Tallquist, M. D., and P. Soriano. 2000. Epiblast-restricted Cre expression in MORE mice: a tool to distinguish embryonic vs. extra-embryonic gene function. *Genesis* 26: 113-115.
3. Stetson, D. B., and R. Medzhitov. 2006. Recognition of cytosolic DNA activates an IRF3-dependent innate immune response. *Immunity* 24: 93-103.
4. Saito, T., D. M. Owen, F. Jiang, J. Marcotrigiano, and M. Gale, Jr. 2008. Innate immunity induced by composition-dependent RIG-I recognition of hepatitis C virus RNA. *Nature* 454: 523-527.