Appendix 1. Baseline Characteristics of Included and Excluded Parturients

	Included (%)	Excluded* (%)
	n=690,742	n=1,714,553
Maternal Age		
<20	44,277 (6.4)	138584 (8.1)
20-34	496,969 (71.9)	1263738 (73.7)
35-39	119,738 (17.3)	251224 (14.7)
≥40	29,758 (4.3)	61007 (3.6)
Race		
Non-Hispanic white	371,615 (53.8)	795199 (46.4)
Non-Hispanic black	99,624 (14.4)	281527 (16.4)
Hispanic	113,495 (16.4)	223342 (13)
Asian or Pacific Islander	39,644 (5.7)	90254 (5.3)
Other/missing	66,363 (9.6)	324231 (18.9)
Annual delivery volume		
≤1000	53,771 (7.8)	277,960 (16.2)
1001-2000	131,438 (19.0)	428,812 (25.0)
2001-4000	323,468 (46.8)	719,854 (42.0)
>4000	182,065 (26.4)	287,927 (16.8)
Antepartum conditions		
Placental abruption	6,174 (0.9)	14,791 (0.9)
Placenta previa	4,624 (0.7)	10,367 (0.6)

Myhre JM, Shilkrut A, Kuklina EV, Callaghan WM, Creanga AA, Kaminsky S. Massive blood transfusion during hospitalization for delivery in New York State, 1998-2007. Obstet Gynecol 2013;123.

The authors provided this information as a supplement to their article.

Previous cesarean delivery	95,802 (13.9)	225,285 (13.1)
Intrauterine fetal demise	4,310 (0.6)	11,385 (0.7)
Risk factors for atony		
Chorioamnionitis	11,517 (1.7)	28,843 (1.7)
Multiple gestation	16,355 (2.4)	31,177 (1.8)
Leiomyomas	8,876 (1.3)	19,080 (1.1)
Severe preeclampsia	6,440 (0.9)	16,813 (1.0)
Induction of labor	86,967 (12.6)	190,937 (11.1)
Cesarean delivery	207,899 (30.1)	473,361 (27.6)
Hemorrhagic complications		
Uterine atony	11,344 (1.6)	28,959 (1.7)
Retained placenta with	1,858 (0.3)	4,637 (0.3)
postpartum hemorrhage		

^{*}All deliveries from hospitals that did not record a single unit of blood transfused during the year of delivery were excluded from further analysis.

Myhre JM, Shilkrut A, Kuklina EV, Callaghan WM, Creanga AA, Kaminsky S. Massive blood transfusion during hospitalization for delivery in New York State, 1998-2007. Obstet Gynecol 2013;123.

The authors provided this information as a supplement to their article.