

Table S1. Procedures and comorbidities of the cohort.

	N	(%)
	66,105	(100)
<i>Preoperative albumin</i>		
<3.5	2148	(3.3)
3.5-4	9234	(14.0)
>4	13,086	(19.8)
Unknown	41,637	(63.0)
<i>Indication for surgery</i>		
Leiomyoma	17,746	(26.9)
Endometriosis	3619	(5.5)
Abnormal bleeding	12,955	(19.6)
Benign neoplasm	2535	(3.8)
Pelvic organ prolapse	7893	(11.9)
Uterine cancer	6683	(10.1)
Cervical cancer	869	(1.3)
Ovarian cancer	1626	(2.5)
<i>Concomitant procedures</i>		
Anterior colporrhaphy	4045	(6.1)
Posterior colporrhaphy	3405	(5.2)
Incontinence repair	3700	(5.6)
Sacral colpopexy	1869	(2.8)
Sacrospinous fixation	774	(1.2)
Lymphadenectomy	4045	(6.1)
<i>Preoperative conditions</i>		
Non insulin dependent diabetes mellitus	3439	(5.2)
Insulin dependent diabetes mellitus	1392	(2.1)
Tobacco use	11,825	(17.9)
Chronic obstructive pulmonary disease	850	(1.3)
Congestive heart failure	63	(0.1)
Hypertension	19,793	(29.9)
Metastatic cancer	838	(1.3)
Corticosteroid use	732	(1.1)
Weight loss	301	(0.5)
Bleeding disorder	744	(1.1)
Preoperative transfusion	488	(0.7)
<i>Postoperative conditions</i>		
Reoperation	1045	(1.6)
Superficial surgical site infection	1051	(1.6)
Deep surgical site infection	219	(0.3)
Organ space surgical site infection	617	(0.9)
Any wound infection	1862	(2.8)
Dehiscence	235	(0.4)
Pneumonia	207	(0.3)
Pulmonary embolism	217	(0.3)
Deep vein thrombosis	171	(0.3)
Urinary tract infection	1842	(2.8)
Transfusion	2742	(4.2)
Sepsis	439	(0.7)
Shock	78	(0.1)
Myocardial infarction	42	(0.1)
Acute renal failure	30	(0.1)
Readmission	1719	(2.6)

Procedures >1% included