

S1 Table. Primer sequences used for quantitative RT-PCR

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
rat TGF β 1 (<i>Tgfb1</i>)	ACCTGCAAGACCATCGACATG	CGAGCCTTAGTTTGGACAGGAT
rat CTGF (<i>Ctgf</i>)	AAGACCTGTGGGATGGGC	TGGTGCAGCCAGAAAGCTC
rat Collagen 1 (<i>Col1a1</i>)	CAACCTCAAGAAGTCCCTGC	AGGTGAATCGACTGTTGCCT
rat Collagen 3 (<i>Col3a1</i>)	CACCCCTCTCTTATTTTGGCAC	AGACTCATAGGACTGACCAAGGTAGTT
rat TNF α (<i>Tnf</i>)	CCCAGACCCTCACACTCAGAT	TTGTCCCTTGAAGAGAACCTG
rat MCP1 (<i>Ccl2</i>)	ATGCAGTTAATGCCCCACTC	TTCCTTATTGGGGTCAGCAC
rat Nox2 (<i>Cybb</i>)	ACTTCTTGGGTCAGCACTGG	GTTCCCTGTCCAGTTGTCTTCG
rat Nox4 (<i>Nox4</i>)	TAGCTGCCCACTTGGTGAACG	TGTAACCATGAGGAACAATACCACC
rat ANP (<i>Nppa</i>)	ATCACCAAGGGCTTCTTCCT	TGTTGGACACCGCACTGTAT
rat BNP (<i>Nppb</i>)	ACAATCCACGATGCAGAAGCT	GGGCCTTGGTCCTTTGAGA
rat GAPDH (<i>Gapdh</i>)	ATGATTCTACCCACGGCAAG	CTGGAAGATGGTGATGGGTT
rat β -actin (<i>Actb</i>)	GGGAAATCGTGCGTGACATT	GCGGCAGTGGCCATCTC