⇒

Published online before print June 15, 2014, doi: 10.3899/jrheum.131227 The Journal of Rheumatology July 1, 2014 vol. 41 no. 7 1379-1384

www.jrheum.org



Published online before print June 15, 2014, doi: 10.3899/jrheum.131227 The Journal of Rheumatology July 1, 2014 vol. 41 no. 7 1379–1384

Behçet Syndrome Manifestations and Activity in the United States versus Turkey — A Cross-sectional Cohort Comparison

APPENDIX 1.

Definitions of Organ Involvement in Behçet syndrome.

- 1. Oral ulcer the presence of an oral ulcer not known to be infectious in origin
- 2. Genital ulcer the presence of a genital ulcer not known to be infectious in origin
- 3. Skin disease acneiform lesions, pustular skin disease, or nodular disease
- 4. Arthralgia the presence of joint pain with or without arthritis on exam
- 5. Eye disease uveitis or retinal vasculitis
- 6. Gastrointestinal disease documented presence of a gastrointestinal ulcer on endoscopy. The usage of nonsteroidal antiinflammatory (NSAID) medications was not an exclusion to endoscopy; however, patients were asked to hold NSAID for a minimum of 4 weeks at the US National Institutes of Health and 3 months at the University of Istanbul
- 7. Neurologic disease dural sinus thrombosis or white matter changes on radiographic studies in the setting of clinical neurologic symptoms other than headaches
- 8. Thrombosis deep vein or arterial thrombosis on exam or radiographic studies