

**Table e-1:** Events rates for dual antiplatelet vs aspirin alone

	<b>Dual therapy (aspirin and clopidogrel)</b>	<b>Single therapy (aspirin)</b>	<b>p Value<sup>a</sup></b>
<b>Recurrent ischemic stroke (n=240)</b>	n, events/year (95% CI)	n, events/year (95% CI)	
Neither	30, 1.3 (0.90, 1.9)	42, 2.0 (1.4, 2.6)	
METS only	22, 1.6 (1.0, 2.4)	25, 1.8 (1.2, 2.6)	
DM only	8, 2.7 (1.2, 4.9)	6, 1.9 (0.71, 3.8)	
Both	46, 2.9 (2.1, 3.8)	61, 3.6 (2.7, 4.5)	0.64
<b>Incident MI (n=74)</b>			
Neither	9, 0.39 (0.18, 0.68)	7, 0.31 (0.13, 0.58)	
METS only	6, 0.43 (0.16, 0.84)	12, 0.84 (0.44, 1.4)	
DM only	4, 1.3 (0.35, 2.8)	3, 0.96 (0.20, 2.3)	
Both	14, 0.84 (0.46, 1.3)	19, 1.3 (0.62, 1.6)	0.53

Abbreviations: METS = metabolic syndrome; DM = diabetes mellitus; MI = myocardial infarct.

<sup>a</sup>Calculated by Wald test for overall differences.

**Table e-2:** Events rates for of blood pressure target

	<b>Higher (SBP 130-139mmHg)</b>	<b>Lower (SBP&lt;130mmHg)</b>	<b>p Value<sup>a</sup></b>
<b>Recurrent ischemic stroke (n=240)</b>	n, events/year (95% CI)	n, events/year (95% CI)	
Neither	34, 1.6 (1.1, 2.2)	38, 1.7 (1.2, 2.3)	
METS only	29, 2.1 (1.4, 2.9)	18, 1.3 (0.80, 2.0)	
DM only	8, 2.9 (1.2, 5.1)	6, 1.9 (0.68, 3.6)	
Both	57, 3.5 (2.7, 4.5)	50, 3.0 (2.3, 3.9)	0.56
<b>Incident MI (n=74)</b>			
Neither	8, 0.36 (0.16, 0.65)	8, 0.34 (0.15, 0.62)	
METS only	11, 0.76 (0.38, 1.3)	7, 0.51 (0.21, 0.96)	
DM only	5, 1.7 (0.55, 3.5)	2, 0.60 (0.073, 1.7)	
Both	15, 1.0 (0.61, 1.5)	18, 1.0 (0.61, 1.5)	0.55

Abbreviations: METS = metabolic syndrome; DM = diabetes mellitus; MI = myocardial infarct.

<sup>a</sup>Calculated by Wald test for overall differences.