

SUPPLEMENTARY TABLE S4. REPRODUCIBILITY OF TRACT-DERIVED FEATURE (SD, PEAK HEIGHT, AND SKEWNESS) OF GFA AND MTR BETWEEN TP1 AND TP2 IN HEALTHY CONTROLS. PERCENTAGE DIFFERENCES (RELATIVE CHANGE [%]) IN TRACT-DERIVED FEATURE (SD, PEAK HEIGHT, AND SKEWNESS) AND MEAN GFA/MTR BETWEEN TP1 AND TP2 IN HEALTHY CONTROLS

		<i>Controls (n = 10. tp1 vs. tp2). Relative change (%). Positive (negative): increase (decrease)</i>											
		<i>Intrahemispheric</i>											
<i>ROI₁-ROI₂</i>		<i>MI-SMA</i>		<i>PMd-MI</i>		<i>PMd-SMA</i>		<i>PMd-PMv</i>		<i>PMv-SMA</i>			
<i>Features</i>	<i>GFA/MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>
Mean		1 ± 3	0 ± 7	2 ± 3	1 ± 5	-10 ± 37	-7 ± 33	1 ± 3	0 ± 5	1 ± 3	0 ± 5	2 ± 4	-1 ± 6
SD		-2 ± 13	-4 ± 25	1 ± 12	5 ± 28	7 ± 32	15 ± 40	1 ± 20	-2 ± 25	1 ± 20	-2 ± 25	-13 ± 36	4 ± 23
Peak Height		32 ± 23	37 ± 95	2 ± 233	-75 ± 305	-42 ± 181	360 ± 653	81 ± 106	750 ± 1401	81 ± 106	750 ± 1401	233 ± 418	-991 ± 3506
Skewness		8 ± 19	11 ± 21	-8 ± 31	-27 ± 45	-40 ± 94	-244 ± 690	-1 ± 30	-6 ± 36	-1 ± 30	-6 ± 36	6 ± 35	4 ± 29
		<i>Intrahemispheric</i>											
<i>ROI₁-ROI₂</i>		<i>CC-m-PMd</i>		<i>CC-m-MI</i>		<i>CC-m-SMA</i>							
<i>Features</i>	<i>GFA/MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>	<i>GFA</i>	<i>MTR</i>
Mean		-1 ± 9	-1 ± 6	-4 ± 7	0 ± 4	-1 ± 4	0 ± 5	0 ± 4	1 ± 5	0 ± 4	1 ± 5	0 ± 4	1 ± 5
SD		1 ± 12	-7 ± 56	-5 ± 21	-5 ± 62	-4 ± 15	6 ± 48	11 ± 29	-5 ± 29	11 ± 29	-5 ± 29	11 ± 29	-5 ± 29
Peak Height		162 ± 297	-27 ± 178	124 ± 158	-76 ± 285	120 ± 202	-150 ± 723	48 ± 107	877 ± 2758	48 ± 107	877 ± 2758	48 ± 107	877 ± 2758
Skewness		-13 ± 24	-32 ± 105	8 ± 26	-34 ± 107	13 ± 11	-42 ± 92	-33 ± 70	-13 ± 21	-33 ± 70	-13 ± 21	-33 ± 70	-13 ± 21

Positive (negative) value denotes an increase (decrease) in the GFA/MTR measure in control relative to the first time point (tp1).