SUPPLEMENT

eTable 1: Composite Antidepressant Score CAD. Guidelines for the CAD Scores to Measure Intensity of Antidepressant Medication Treatment (Daily Dosages) – Revised for Older Adults.

eTable 2: Treatment Satisfaction Scores. LS means of Patient and Caregiver Treatment Satisfaction Scores in PATH and ST at 4, 8, 12 Weeks of Treatment, Based on Mixed-Effects Models Analyses.

eTable 1: Composite Antidepressant Score CAD. Guidelines for the CAD Scores to Measure Intensity of Antidepressant Medication Treatment (Daily Dosages) – Revised for Older Adults.

CAD Score	Nortriptyline Pamelor	Sertraline Zoloft	Bupropion Wellbutrin	Fluoxetine Prozac Paroxetine Paxil Citalopram Celexa	Venlafaxine Effexor	Escitalopram Lexapro
0	None	None	None	None	None	None
1	<30mg	<25mg	<100mg	<10mg	<50mg	<5mg
2	30-49mg	25-49mg	100-199mg	10-19mg	50-75mg	5-9mg
3	50-80mg	50-100mg	200-300mg	20-30mg	76-150mg	10-15mg
4	>80mg	>100mg	>300mg	>30mg	>150mg	>15mg

CAD Score	Mirtazapine Remeron	Fluroxamine Luvox	Nefazodone Serzone	Duloxetine Cymbalta
0	None	None	None	None
1	<10mg	<50mg	<150mg	<20mg
2	10-15mg	50-99mg	150-250mg	21-40mg
3	16-30mg	100-200mg	251-350mg	41-60mg
4	>30mg	>200mg	>350mg	61-120mg

eTable 2: Treatment Satisfaction Scores. LS means of Patient and Caregiver Treatment Satisfaction Scores in PATH and ST at 4, 8, 12 Weeks of Treatment, Based on Mixed-Effects Models Analyses.

modele / maryeee.	PAT PATIENT Mean (SE)	H (N=37) CAREGIVER Mean (SE)	ST (I PATIENT Mean (SE)	N=37) CAREGIVER Mean (SE)					
To What Extent Did Service Meet Patient's Needs?*§									
Week 4	3.01 (0.16)	3.05 (0.25)	3.02 (0.15)	3.02 (0.27)					
Week 8	3.12 (0.13)	3.37 (0.21)	3.08 (0.13)	3.07 (0.24)					
Week 12	3.23 (0.15)	3.68 (0.24)	3.13 (0.14)	3.13 (0.28)					
What Was the Overall Satisfaction With Service? *§									
Week 4	3.48 (0.14)	3.51 (0.18)	3.47 (0.13)	3.37 (0.20)					
Week 8	3.46 (0.11)	3.59 (0.16)	3.51 (0.11)	3.39 (0.18)					
Week 12	3.44 (0.13)	3.67 (0.18)	3.56 (0.12)	3.41 (0.21)					
Would Patient Come Back To The Program ? *§									
Week 4	3.70 (0.13)	3.78 (0.16)	3.64 (0.12)	3.41 (0.17)					
Week 8	3.67 (0.11)	3.78 (0.15)	3.63 (0.11)	3.49 (0.16)					
Week 12	3.65 (0.12)	3.79 (0.16)	3.63 (0.11)	3.57 (0.18)					

*Client Satisfaction Questionnaire:

^{1:} To What Extent Did Service Meet Patient's Needs? (Score range 1-4: 4=Almost all needs have been met; 3=Most needs met; 2=Only a few needs met; 1=No needs met);

^{2:} Overall Satisfaction With Service (Scores range 1-4; 4=Very satisfied; 3=Mostly satisfied; 2=Indifferent or mildly dissatisfied; 1=Quite dissatisfied);

^{3:} Would You Come Back To The Program? (Scores range 1-4; 4=Yes, definitely; 3=Yes, I think so; 2=No, I don't think so; 1=No, definitely not).

[§] Mixed-effects models analysis revealed that there were no significant differences in slopes between the two treatments (treatment group x time interaction was not significant) on each CSQ item. Further, there were no significant differences between the two treatments at each time point, as shown by post-hoc analysis with Bonferroni adjustment for multiple comparisons.