

Title page:

Inhibition of the Rho/Rho kinase pathway prevents lipopolysaccharide-induced hyperalgesia and the release of TNF- α and IL-1 β in the mouse spinal cord

Author: Cunjin Wang^{1#}, Siyuan Song^{2#}, Yang Zhang¹, Yali Ge¹, Xiangzhi Fang¹, Tianfeng Huang¹, Jin Du¹ and Ju Gao^{1*}

1 Clinical Medical College of Yangzhou University & Department of Anesthesiology, Subei People's Hospital of Jiangsu Province, Yangzhou, China

2 Jiangsu Key Laboratory of Anesthesiology & Jiangsu Key Laboratory of Anesthesia and Analgesia Application Technology, Xuzhou Medical College, Xuzhou, China

These authors contributed equally to this work.

*Corresponding Author: Ju Gao. Address: No.98 Nan Tong Western Road, Yangzhou, Jiangsu, 225001, China. E-mail: gaoju_003@163.com. Tel: +86-0514-87373650.

Supplementary figure legend

Dose-related effects of C3 exoenzyme (A), (B) and Y27632 (C), (D) on the pain behavior in LPS-treated mice. C3 exoenzyme, Y27632 or saline was administered intrathecally before LPS. The mechanical hyperalgesia and thermal hyperalgesia were measured 3h after the treatment with LPS. One-way ANOVA, $*P < 0.05$ compared with saline control, $n = 8$ mice in each group.

