

**Efficacy and HIV drug resistance profile of second-line ART among patients receiving long-term first-line regimens in rural China**

Jing Wang<sup>1,2#</sup>, Zhe Wang<sup>3#</sup>, Jia Liu<sup>3#</sup>, Yanchao Yue<sup>4</sup>, Shimei Yang<sup>5</sup>, Huimin Huang<sup>6</sup>, Cui He<sup>1</sup>, Lingjie Liao<sup>1</sup>, Hui Xing<sup>1\*</sup>, Yuhua Ruan<sup>1</sup>, Yiming Shao<sup>1\*</sup>

**Supplementary Table S1. Prevalence of HIV drug resistance mutations before and at 6 and 12 months after switching drugs**

	Before switch regimen N=90 (%)	6 month after switch regimen N=81 (%)	12 month after switch regimen N=82 (%)
NNRTIs	59(65.5)	20(24.7)	22(26.8)
K101EN	4(4.4)	1(1.2)	0
K103NS	35(38.9)	13(16.0)	16(19.5)
Y181CY	20(22.2)	11(13.6)	10(12.2)
Y188LFY	6(6.7)	2(2.5)	1(1.2)
G190A	13(14.4)	1(1.2)	2(2.4)
NRTIs	48(53.3)	16(19.7)	15(18.3)
M41L	19(21.1)	6(7.4)	8(9.7)
K65R	1(1.1)	0	0
L74V	1(1.1)	1(1.2)	1(1.2)
D67NTG	18(20.0)	6(7.4)	7(8.5)
K70R	14(15.5)	6(7.4)	6(7.3)
T69NDAT	8(8.9)	3(3.7)	4(4.9)
M184VI	44(48.9)	13(16.0)	13(15.8)
T215YSF	31(34.4)	12(14.8)	10(12.2)
L210W	8(8.9)	3(3.7)	3(3.6)
PIs	1(1.1)	0	0
I47V	1(1.1)	0	0