

S3 Table. Anthropometry by visceral fat missing status in the MESA body composition ancillary study

Variable	Non-Missing Mean (SD)	Missing Visceral Fat Mean (SD)	Imputed Visceral Fat Mean (SD)	P-value: Non-Missing vs. Missing
N	1898	24	25	
Visceral fat (cm ²)	149.6 (1.61)	NA	258.8 (8.13)	
Age (years)	62.0 (0.22)	64.8 (1.91)	64.3 (2.13)	<0.001
Female (%)	50.1 (1.15)	16.7 (7.78)	32.0 (9.52)	<0.001
Height (cm)	166.3 (0.23)	170.7 (1.51)	170.7 (2.82)	<0.001
Weight (kg)	77.6 (0.38)	101.2 (4.02)	97.3 (2.79)	<0.001
BMI (kg/m ²)	28.0 (0.12)	34.6 (1.14)	33.6 (0.98)	<0.001
Waist circumference (cm)	97.8 (0.32)	118.5 (2.70)	113.6 (1.92)	<0.001
Hip Circumference (cm)	104.1 (0.25)	115.4 (2.12)	114.3 (2.06)	<0.001
Waist to Hip Ratio	0.94 (0.002)	1.03 (0.013)	1.00 (0.009)	<0.001
Waist to Height Ratio	0.59 (0.002)	0.69 (0.015)	0.67 (0.019)	<0.001