

Table e-1: chi-square test

Variable	Level	Completed recurrent stroke survey N (%)	Did not complete recurrent stroke survey N (%)	Total	P-value (based chi-square)
Total		329	160	489	
Gender	Male	173 (53%)	85 (53%)	258	0.91
	Female	156 (47%)	75 (47%)	231	
Race	White non-hispanic	263 (80%)	132 (83%)	395	0.75
	Black	19 (6%)	6 (4%)	25	
	Other	44 (13%)	20 (13%)	64	
	Unknown	3 (<1%)	2 (1%)	5	
Age at cancer diagnosis, years	0-4	116 (35%)	49 (31%)	165	0.58
	5-9	76 (23%)	41 (26%)	117	
	10-14	69 (21%)	40 (25%)	109	
	15-20	68 (21%)	30 (19%)	98	
Age at first stroke, years	0-17	142 (43%)	69 (43%)	211	0.20
	18-29	66 (20%)	28 (18%)	94	
	30-39	70 (21%)	30 (19%)	100	
	≥40	32 (10%)	14 (9%)	46	
	Unknown	19 (6%)	19 (12%)	38	
Age at last followup or death, years	0-17	22 (7%)	13 (8%)	35	0.48
	18-29	91 (28%)	51 (32%)	142	
	30-39	122 (37%)	60 (38%)	182	
	≥40	94 (29%)	36 (23%)	130	
Cancer diagnosis	Brain tumor	140 (43%)	61 (38%)	201	0.68
	Leukemia	90 (27%)	51 (32%)	141	
	Hodgkins	35 (11%)	15 (9%)	50	
	Other	64 (19%)	33 (21%)	97	
Cranial radiation therapy maximum dose*	None	55 (18%)	25 (19%)	80	0.15
	1-29 Gy	55 (18%)	32 (24%)	87	
	30-49 Gy	30 (10%)	16 (12%)	46	
	50+ Gy	97 (31%)	27 (21%)	124	
	Indirect	52 (17%)	18 (14%)	70	
	Unknown	21 (8%)	13 (10%)	82	

- Among those with complete medical records abstraction