Supplementary Online Content

Ernst KF, Hall DE, Schmid KK, et al. Surgical palliative care consultations over time in relationship to system-wide frailty screening. JAMA Surg. Published online September 10, 2014. doi:10.1001/jamasurg.2014.1393.

eAppendix. Risk analysis index

This supplementary material has been provided by the authors to give readers additional information about their work.

eAppendix. Risk Analysis Index

Last Name			Last four SSN					
Date this Fo	orm completed_		·					
Date of Surgery-Anticipated								
Proposed O	peration							
	AND CANCI							
1. Sex				Female = 0)/ Male =			
5								
10A- Age and cancer Score	Score	Score	2.	Age				
Age	without cancer	with cancer	(Excluding skin cane	er, except for melanoma)				
< + 69	2	20	_ (Environing simil value	Score with Cancer				
70 – 74	3	19						
75 – 79	4	18	_	OR				
80- 84	5	17	_	Score without cancer				
85 – 89	6	15	4					
90 -94	7	14	4					
95 -99 100+	9	14	4					
100+	9	13	_					
MEDIA	CAL CO MOI	DIDITIES	y.					
	CAL CO-MOI				Na/Vas			
	•	ntentional w	weight loss in the past 3 n	nonths (>10 lbs)?	No/Yes			
	5							
3. Renal failure?								
No/Yes 6								
4. Chronic/congestive heart failure?								
No/Yes 4								
	_	_4						
5. Po	or appetite?							
	No/Yes_	_4						
6. Sh	ortness of breat							
5. Bii								
	No/Yes_	_o						

COGNITION, RESIDENCE, AND ACTIVITY OF DAILY LIVING:

Cognition and Activities of Daily Living:

7. Do you reside at a setting other than independent living? (listed below)

No/Yes_8__

If YES, circle answer: Skilled Nursing Facility/ Assisted Living/ Nursing Home If YES, admission within in the past 3 months?

No/Yes

8. Have your cognitive skills or status deteriorated over the last 3 months?

No/Yes

9. Activities of Daily Living Chart:

Without cognitive decline

____ (0-16)

____ (-2 to 21)

With cognitive decline

TOTAL SCORE

PERCENT

Mobility/locomotion		Eating		Toilet use		Personal hygiene	
0.	Independent	0.	Independent	0.	Independent	0.	Independent
1.	Supervised	1.	Supervised	1.	Supervised	1.	Supervised
2.	Limited assistance	2.	Limited assistance	2.	Limited assistance	2.	Limited assistance
3.	Extensive assistance	3.	Extensive	3.	Extensive	3.	Extensive
4.	Total dependence	assistance			assistance		assistance
	-	4.	Total dependence	4.	Total dependence	4.	Total dependence

INDEPENDENT = No help or oversight – OR – help or oversight provided only 1 or 2 times in last 7 days SUPERVISED = Oversight, supervision or cueing provided 3 or more times during the last 7 days.

LIMITED ASSISSTANCE= patient highly involved in activity; but received physical help in guided maneuvering of limbs or other non weight bearing assistance 3 or more times in the last 7 days.

EXTENSIVE ASSISSTANCE= while patient performed *part* of activity over last 7 day period, help was provided for the following: weight bearing support OR full staff performance during part of the past 7 days. TOTAL DEPENDENCE= full staff performance during the past 7 days.

TOTAL DEPENDENCE= full staff performance during the past 7 days.

12- ADL POINTS SCORE	ADL POINTS SCORE with COGNITIVE DECLINE
0	ADL score -2
1,2	ADL score -1
3,4	ADL score 0
5,6,7	ADL score +1
8,9	ADL score +2
10,11	ADL score +3
12,13	ADL score +4
14,15,16	ADL score +5

Points	=	Points =	Points =
percent		percent	percent
0-5	=	26 – 30 =	56- 60 =
4%		27%	89%
6- 10	=	31 – 35 =	61 65 =
4%		36%	90%
11-15	=	36 – 40 =	66- 70 =
7%		47%	93%
16-20	=	41 – 45 =	71 -75 =
11%		58%	100 %
21- 25	=	46- 50 =	
17%		69%	