

**Table S1. Demographic and clinical characteristics of HIV-infected African-American women in WIHS, stratified by inclusion/exclusion status for APOL1 substudy**

<b>Characteristic</b>	<b>Included (N = 431)</b>	<b>Excluded (N = 364)</b>	<b>P-value</b>
<b>Baseline Age (y)</b>	41 (36-46)	42 (36-47)	0.55
<b>Hypertension</b>	31%	34%	0.37
<b>Diabetes mellitus</b>	10%	19%	0.0006
<b>Hepatitis C virus infection</b>	35%	37%	0.56
<b>Current CD4+ cell count (cells/mm<sup>3</sup>)</b>	391 (246-583)	330 (179-539)	0.0011
<b>History of AIDS</b>	48%	48%	0.93
<b>HIV RNA (copies/mL)</b>			
<b>&lt;=80</b>	28%	20%	0.087
<b>81-1999</b>	23%	27%	
<b>2000-9999</b>	16%	17%	
<b>&gt;=10000</b>	32%	36%	
<b>Current HAART use</b>	59%	32%	<.0001

Data are presented as median (interquartile range) or numbers (percent). P-values from chi-square and Kruskal-Wallis tests for categorical and continuous variables, respectively.

Abbreviations: HAART, highly active antiretroviral therapy.