$\label{eq:instructions} INSTRUCTIONS \\ to complete the DM1-ACTIV^{C@}$

DM1-ACTIV^{C©} General aspects

- This 25 item questionnaire covers a wide range of daily and social activities and represents the preliminary 146 items based on the WHO-ICF classification of activities and participation from which it was derived.
- The DM1-ACTIV^{C©} captures personal ability to execute a task where the patient is requested to determine whether he/she is able to complete such a task independently.

		Impossible to perform	Able to perform, but	Able to perform, without
			with difficulty	difficulty
	Item	(0)	(1)	(2)
	Are you able to			
1	eat soup?			
2	visit family or friends?			
3	care for your hair and body?			
4	dress your lower body?			
5	wash your upper body?			
6	take a shower?			
7	wash your lower body?			
8	get out of bed?			
9	move a chair?			
10	do the dusting/cleaning?			
11	do the shopping?			
12	tie the laces of your shoes?			
13	catch an object (e.g. a ball)?			
14	use dustpan and brush?			
15	empty dustbin?			
16	make up your bed?			
17	vacuum clean?			
18	serve coffee/tea on a tray?			
19	dance?			
20	stand up from squatting position?			
21	stand on one leg?			
22	walk uphill?			
23	walk 3 flights of stairs?			
24	carry and put down heavy object (10 kg)?			
25	run?			

DM1-ACTIV^{C©} General aspects

- Patients should complete the questionnaire after having received written and verbal instructions.
- Make sure that the patient is able to complete the questionnaire in a comfortable and quiet room.
- The patient should be comfortably sitting.
- Encourage the patient to complete the questions by themselves.
 However, a partner, relative or friend may be of assistence if the patient is unsure or has communication problems.
- It takes approximately 3 minutes to complete the questionnaire.

DM1-ACTIV^{C©} Instructions-I

- Make sure that the answers given are due to the impact of myotonic dystrophy type I, rather than another concomitant disease (e.g. arthritis).
- Ask the patient to rate the tasks as either "Impossible to perform", "Able to perform, but with difficulty" or "Able to perform, without difficulty".
- If it is the first time for the patient completing the DM1-ACTIV^{C©}, go through each question with the patient and ensure the patient that the interpretation of each question is left to his/her best personal judgement rather than a fixed way to interpret each question.

DM1-ACTIV^{C©} Instructions-II

- Instruct the patient that there are no right or wrong answers. Encourage the patient to choose the response that best describes his/her situation.
- A partner, family or a carer may help in interpreting the tasks in the way that reflects the situation of the patient as accurate as possible.
- For example: "Able to do the shopping?" The patient may experience difficulties at various levels, which may lead to problems executing such a task. This may also differ from patient to patient.
- Problems executing this task may be due to physical impairment such as weakness leading to mobility problems, transportation limitations, financial handling problems, executive cognitive dysfunctions.

DM1-ACTIV^{C©} Instructions-III

- In case of any doubt or addressing a task that is not relevant to the patient (e.g. "I never do vacuum cleaning"), the patient is requested to choose an answer to the best of his/her personal judgment of being able or not to perform such a task, with or without difficulty.
- If still unsure and if possible, you may observe the patient performing the task, or miming the task.
- When using a special device (for example adapted cutlery (spoon), special cooking utensils, cane, crutch, or walker) to complete a task, the patient is requested to score "Able to perform, but with difficulty".
- However, if the patient is not able to execute a task despite using special devices, the patient should score "Impossible to perform".

DM1-ACTIV^{C©} Instructions-IV

- When some assistance is needed (for example support by a caregiver), to complete a task, the patient is requested to score "Able to perform, but with difficulty".
- However, if the patient is not able to contribute in completing a task, the patient should score "Impossible to perform".