

Supplementary Table 2. Patients characteristics according to lymphadenectomy

Parameter	No lymphadenectomy (n=89)	Pelvic +/-aortic lymphadenectomy (n=165)	p-value*
Age (yr)			0.253
≤65	51 (57.3)	103 (62.4)	
>65	38 (42.7)	62 (37.6)	
Stage			<0.001
IB	28 (31.5)	33 (20.0)	
II	33 (37.1)	40 (24.2)	
IIIA	26 (29.2)	28 (17.0)	
IIIB	2 (2.3)	6 (3.6)	
IIIC	0 (0)	58 (35.2)	
Myometrial invasion (%)			0.507
No	0	0	
≤50	26 (29.2)	47 (28.5)	
>50	63 (70.8)	118 (71.5)	
Grade			0.134
1 and 2	37 (41.6)	82 (49.7)	
3	52 (58.4)	83 (50.3)	
LVSI			0.213
Negative	62 (69.7)	104 (63.8)	
Positive	27 (30.3)	59 (36.2)	
Adjuvant therapy			0.224
No	20 (22.5)	35 (21.2)	
EBRT	41 (46.1)	57 (34.6)	
Chemo	17 (19.1)	42 (25.5)	
Sequential chemo-EBRT	11 (12.4)	31 (18.8)	

Values are presented as number (%).

EBRT, external beam radiotherapy; LVSI, lymphovascular space involvement.

*Fisher exact test or Student t-test.